

NCHS_HIW 2.2 Data Release 2

ShortDescription	FullDescription	National	State	County	Hospital Referral Region	HIW ID
Abstinence discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about abstinence before they were 18	Yes				810
Abstinence discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about abstinence before they were 18	Yes				811
Abstinence instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on abstinence before they were 18 years	Yes				802
Abstinence instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on abstinence before they were 18 years	Yes				803
Activity limitation due to arthritis: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who experience a limitation in activity due to arthritis	Yes				419
Activity limitation due to chronic back conditions (per 1,000)	Activity limitation due to chronic back conditions	Yes				428
Activity limitations: adults 45+ years with COPD (percent)	Percent of adults aged 45 years and over with chronic obstructive pulmonary disease (COPD) who have activity limitations	Yes				1405
Activity limitations: persons with asthma (percent)	Percent of persons with current asthma who have activity limitations	Yes				1404
Acute chemical poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that cause acute chemical poisoning	Yes				718
Acute hospital readmissions (count)	Total inpatient readmissions within 30 days of an acute hospital stay	Yes	Yes	Yes	Yes	278
Acute hospital readmissions (percent)	Inpatient readmissions within 30 days of an acute hospital stay	Yes	Yes	Yes	Yes	279
Added sugar consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from added sugars for persons aged 2 years and over	Yes				1237
ADHD: children (percent)	Percent of children under age 18 years who have attention deficit hyperactivity disorder (ADHD)	Yes				50033
Adolescents having an adult to discuss problems with (percent)	Percent of adolescents who have an adult in their lives with whom they can talk about serious problems	Yes				409
Adolescents' parents attend activities/events (percent)	Percent of parents who attend events and activities in which their adolescents participate	Yes				410
Aerobic physical activity meeting guidelines: adults (percent)	Percent of adults who engage in aerobic physical activity of at least moderate intensity for at least 150 minutes per week	Yes				1319
Aerobic physical activity meeting guidelines: teens (percent)	Percent of adolescents who meet current Federal physical activity guidelines for Aerobic physical activity	Yes	Yes			1323
Aerobic physical activity meeting high guidelines: adults (percent)	Percent of adults who engage in aerobic physical activity of at least moderate intensity for more than 150 minutes per week	Yes				1320
Aerobic physical and muscle-strengthening activity: adults (percent)	Percent of adults who meet the objectives for aerobic physical activity and for muscle-strengthening activities	Yes				1322
African American Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who are African American	Yes	Yes	Yes	Yes	312
AIDS, perinatally acquired (number)	Number of new cases of perinatally acquired AIDS	Yes				939
AIDS: heterosexual persons 13+ years (number)	Number of new AIDS cases among adolescent and adult heterosexuals	Yes				943
AIDS: injection drug users (number)	Number of new AIDS cases among adolescents and adults who inject drugs	Yes				931
AIDS: men who have sex with men (number)	Number of new AIDS cases among adolescent and adult men who have sex with men	Yes				930
AIDS: persons 13+ years (per 100,000)	Rate of new AIDS cases among adolescents and adults	Yes				929
Air quality management program: schools (percent)	Percent of the Nation's elementary, middle, and high schools that engage in practices that promote a healthy environment	Yes				683
Airborne toxins from area sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - area sources	Yes				671
Airborne toxins from major sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - major sources	Yes				672
Airborne toxins from mobile sources (number)	Air toxic emissions to decrease the risk of adverse health effects caused by airborne toxics - mobile sources	Yes				670
Alcohol abstinence, preconception (percent)	Percent of women delivering a live birth who did not drink alcohol prior to pregnancy	Yes				1181
Alcohol abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Alcohol	Yes				1142
Alcohol and/or illicit drug treatment receipt (percent)	Percent of persons who need alcohol and/or illicit drug treatment and received specialty treatment for alcohol and/or illicit drug use	Yes				1455
Alcohol Screening and Brief Intervention: primary care & trauma centers (number)	Number of Level I and Level II trauma centers and primary care settings that implement evidence-based alcohol screening and brief intervention	Yes				1460
Alcohol sold (gallons)	Average annual alcohol consumption rate	Yes				1439
Alcohol treatment receipt (percent)	Percent of persons who need alcohol and/or illicit drug treatment and received specialty treatment for alcohol and/or illicit drug use	Yes				1456
Alcohol, never use: at risk teens (percent)	Percent of at risk adolescents aged 12 to 17 years who, in the past year, refrained from using alcohol for at least 12 months	Yes				1428
Alcohol, never use: high school seniors (percent)	Percent of high school seniors never using substances-Alcoholic beverages	Yes				1430
Alcohol-attributable deaths (average annual number)	Number of deaths attributable to alcohol	Yes				1467
Alcohol-impaired driving deaths (per 100,000,000 vehicle miles traveled)	Alcohol-impaired driving (.08+ blood alcohol content [BAC]) fatality rate	Yes				1466
Allergic reactions to food: adults (percent)	Percent of adults with a physician-diagnosed food allergy with a severe allergic reactions to food	Yes				820
Alzheimer's disease deaths (per 100,000)	Alzheimer's disease death rate per 100,000	Yes	Yes	Yes	Yes	50015
Alzheimer's Disease, Related Disorders, or Senile Dementia Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have Alzheimer's Disease, Related Disorders, or Senile Dementia	Yes	Yes	Yes	Yes	50016
Alzheimer's Disease, Related Disorders, or Senile Dementia Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have Alzheimer's Disease, Related Disorders, or Senile Dementia	Yes	Yes	Yes	Yes	50021
Ambulatory surgery center Medicare users (count)	Medicare fee-for-service beneficiaries who used ambulatory surgery center services	Yes	Yes	Yes	Yes	233
Ambulatory surgery center Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used ambulatory surgery center services	Yes	Yes	Yes	Yes	234
Ambulatory surgical center Medicare service events (per 1,000 beneficiaries)	Rate of ambulatory surgery center service events among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2020

Ambulatory Surgical Centers (number)	Number of Medicare participating Ambulatory Surgical Centers	Yes	Yes	Yes	10033
Amputation among diabetics rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Rate of lower extremity amputation among patients with diabetes amor	Yes	Yes	Yes	2046
Anencephaly (per 100,000 live births)	Rate of anencephaly	Yes			1170
Antibiotic use: persons with the common colds (per 100,000)	Rate of courses of antibiotics prescribed for the sole diagnosis of the common cold	Yes			975
Antibiotic-resistant pneumococcal infections: children <5 years (per 100,000)	Rate of invasive antibiotic-resistant pneumococcal infections among children under age 5 years	Yes			965
Antibiotic-resistant pneumococcal infections: older adults (per 100,000)	Rate of invasive antibiotic-resistant pneumococcal infections among adults aged 65 years and older	Yes			966
Antibiotics for ear infections: children <5 years (per 100)	Rate of physician visits by children less than 5 years of age diagnosed with an ear infection that have a	Yes			974
Antimicrobial resistance to 3+ classes of antimicrobials (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant	Yes			833
Antimicrobial resistance to Ampicillin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant	Yes			832
Antimicrobial resistance to Ceftriaxone (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant	Yes			830
Antimicrobial resistance to Erythromycin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant	Yes			834
Antimicrobial resistance to Gentamicin (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant	Yes			831
Antimicrobial resistance to Nalidixic acid (percent)	Percent of non-typhoidal Salmonella and Campylobacter jejuni isolates from humans that are resistant	Yes			829
Appropriate treatment use: adults with balance/dizziness problems (percent)	Percent of adults who have tried recommended methods for treating their balance or dizziness proble	Yes			765
Appropriate treatment use: adults with tinnitus (percent)	Percent of adults for whom tinnitus is a moderate to severe problem who have tried appropriate treat	Yes			761
Arsenic level (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes			695
Arsenic poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that ca	Yes			716
Arteriovenous fistula use: hemodialysis patients (percent)	Percent of adult hemodialysis patients who use arteriovenous fistulas or have a maturing fistula as the	Yes			471
Arthritis education: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who have had effective, evidence-based arthritis edu	Yes			426
Arthritis Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have arthritis	Yes	Yes	Yes	50017
Arthritis Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have arthritis	Yes	Yes	Yes	50022
Artificial UV tanning: adults (percent)	Percent of adults aged 18 and older who report using artificial sources of ultraviolet light for tanning	Yes			502
Artificial UV tanning: teens (percent)	Percent of adolescents in grades 9 through 12 who report using artificial sources of ultraviolet light for	Yes			501
Aspirin use for secondary prevention (percent)	The percentage of physician office visits to primary care physicians and cardiologists by adult patients	Yes			928
Assaults, work-related (per 10,000)	Rate of work-related assaults	Yes			1311
Assistive listening device use: older adults (per 1,000)	Rate of adults aged 70 years and older with hearing loss who use assistive listening devices	Yes			747
Assistive/adaptive devices use: adults with visual impairment (percent)	Percent of persons aged 18 years and over with visual impairment who use assistive and adaptive dev	Yes			1575
Associate degrees in public health: 2-year colleges (percent)	Percent of 2-year colleges that offer public health or related Associate degrees	Yes			1388
Asthma attack in past 12 months: children (percent)	Percent of children under age 18 years who have had an asthma attack in the past 12 months	Yes			50032
Asthma deaths: adults 35-64 years (per 1,000,000)	Asthma death rate: Adults aged 35 to 64 years old	Yes			1408
Asthma deaths: older adults (per 1,000,000)	Asthma death rate: Adults aged 65 years and over	Yes			1409
Asthma deaths: persons <35 years (per 1,000,000)	Asthma death rate: Children and adults under age 35 years	Yes			1407
Asthma ED visits: children <5 years (per 10,000)	Rate of hospital emergency department visits for asthma: Children under age 5 years	Yes			1401
Asthma ED visits: older adults (per 10,000)	Rate of hospital emergency department visits for asthma: Adults aged 65 years and over	Yes			1403
Asthma ED visits: persons 5-64 years (per 10,000)	Rate of hospital emergency department visits for asthma: Children and adults aged 5 to 64 years	Yes			1402
Asthma exposure risks, advise to reduce: persons with asthma (percent)	Percent of persons with current asthma who have been advised by a health professional to change thii	Yes			1417
Asthma hospitalizations: children <5 years (per 10,000)	Rate of hospitalizations for asthma: Children under age 5 years	Yes			1398
Asthma hospitalizations: older adults (per 10,000)	Rate of hospitalizations for asthma: Adults aged 65 years and over	Yes			1400
Asthma hospitalizations: persons 5-64 years (per 10,000)	Rate of hospitalizations for asthma: Children and adults aged 5 to 64 years	Yes			1399
Asthma in younger adults admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Asthma in younger adults admission rate among Medicare fee-for-serv	Yes	Yes	Yes	2045
Asthma inhalers, receipt of instructions (percent)	Percent of persons with current asthma with prescribed inhalers who receive instruction on their use	Yes			1414
Asthma management plans receipt from health care provider (percent)	Percent of persons with current asthma who receive written asthma management plans from their he:	Yes			1413
Asthma Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have asthma	Yes	Yes	Yes	50007
Asthma Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have asthma	Yes	Yes	Yes	50008
Asthma medication regimen, appropriate (percent)	Percent of persons with current asthma who do not use more than one canister of short-acting inhale	Yes			1416
Asthma patient education receipt (percent)	Percent of persons with current asthma who receive formal patient education	Yes			1412
Asthma signs/symptoms and responses, education receipt (percent)	Percent of persons with current asthma who receive education about appropriate response to an asth	Yes			1415
Asthma surveillance systems: states (number)	Numbers of states, territories, and the District of Columbia with a comprehensive asthma surveillance	Yes			1421

Asthma: children (percent)	Percent of children under age 18 years who have asthma at time of interview	Yes				50031
Atrial fibrillation Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have atrial fibrillation	Yes	Yes	Yes	Yes	285
Atrial fibrillation Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have atrial fibrillation	Yes	Yes	Yes	Yes	286
Autism spectrum disorder screening (percent)	Percent of children (10-35 months of age) who are screened for autism and other developmental delays	Yes				1188
Average age of Medicare beneficiaries (mean)	Average age of Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	308
Average total hierarchical condition category score for Medicare beneficiaries (mean)	Average hierarchical condition category score of Medicare beneficiaries with full fee-for-service coverage	Yes	Yes	Yes	Yes	318
Bacterial pneumonia admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Bacterial pneumonia admission rate among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2043
Basic action difficulty/Complex activity limitation: adults (percent)	Percent of persons 18 years and over who have any basic actions difficulty or complex activity limitation	Yes				50049
Basic actions difficulty: adults (percent)	Percent of adults aged 18 and over with at least one basic actions difficulty	Yes				50042
BDE 47 level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations	Yes				712
Beaches open and safe (percent)	Percent of days that beaches are open and safe for swimming	Yes				674
Beta-HCH level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations	Yes				702
Bicycle helmet laws: states (number)	Number of states and the District of Columbia with laws requiring bicycle helmets for bicycle riders	Yes				1049
Bicycling for transportation (percent)	Percent of trips to work using of alternative modes of transportation - bicycling	Yes				666
Binge drinking abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Binge drinking	Yes				1143
Binge drinking: adults (percent)	Percent of persons engaging in binge drinking during the past month-Adults aged 18 years and older	Yes				1437
Binge drinking: adults (percent) (Source: BRFSS)	Percent of adults 18 years and over that report binge drinking		Yes	Yes		17
Binge drinking: college students (percent)	Percent of students engaging in binge drinking during the past 2 weeks-College students	Yes				1436
Binge drinking: high school seniors (percent)	Percent of students engaging in binge drinking during the past 2 weeks-High school seniors	Yes				1435
Binge drinking: teens (percent)	Percent of persons engaging in binge drinking during the past month-Adolescents aged 12 to 17 years	Yes				1438
Birth control discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about birth control methods before they were 18	Yes				812
Birth control discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about birth control methods before they were 18	Yes				813
Birth control instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on birth control methods before they were 18	Yes				804
Birth control instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on birth control methods before they were 18	Yes				805
Births: females 15-19 years (per 1,000)	Births to females 15-19 years of age per 1,000	Yes	Yes	Yes		22
Births: unmarried women 18-54 years (percent)	Percent of all live births to women ages 18-54 years who report not being married	Yes	Yes	Yes		99
Births: women 40-54 years (percent)	Percent of all live births to women 40-54 years of age	Yes	Yes	Yes		98
Births: women under 18 years (percent)	Percent of all live births to mothers less than 18 years of age	Yes	Yes	Yes		97
Bisphenol A level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations	Yes				709
Blood donations: adults (percent)	Percent of persons who donate blood	Yes				436
Blood folate concentration: reproductive-aged women (percent)	Percent of women of childbearing potential who have lower red blood-cell folate concentrations	Yes				1172
Blood pressure control: adults with hypertension (percent)	Percent of adults with hypertension whose blood pressure is under control	Yes				882
Blood pressure screening: adults (percent)	Percent of adults who have had their blood pressure measured within the preceding 2 years and can still see a health care provider	Yes				883
Blood pressure under control: adults with diabetes (percent)	Percent of adults with diagnosed diabetes whose blood pressure is under control	Yes				560
Body mass index assessment of adults by physicians (percent)	Percent of primary care physicians who regularly assess body mass index (BMI) in their adult patients	Yes				1234
Body mass index assessment of children by physicians (percent)	Percent of primary care physicians who regularly assess body mass index (BMI) for age and sex in their patients	Yes				1235
Breakfast program: schools (percent)	Percent of schools with a school breakfast program	Yes				417
Breast cancer deaths (per 100,000)	Breast cancer death (female) rate	Yes	Yes	Yes		488
Breast cancer Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have breast cancer	Yes	Yes	Yes	Yes	299
Breast cancer Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have breast cancer	Yes	Yes	Yes	Yes	300
Breastfeeding, at 1 year (percent)	Percent of infants who are breastfed-At 1 year	Yes				1149
Breastfeeding, at 6 months (percent)	Percent of infants who are breastfed-At 6 months	Yes				1148
Breastfeeding, ever (percent)	Percent of infants who are breastfed-Ever	Yes				1147
Breastfeeding, exclusively through 3 months (percent)	Percent of infants who are breastfed-Exclusively through 3 months	Yes				1150
Breastfeeding, exclusively through 6 months (percent)	Percent of infants who are breastfed-Exclusively through 6 months	Yes				1151
Broadband access to internet (percent)	Percent of persons with broadband access to the Internet	Yes				858
Bullying: teens (percent)	Percent of bullying among adolescents	Yes	Yes			1102
Cadmium level (µg/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine concentrations	Yes				696

Cadmium poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that ca	Yes				717
Campylobacter species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Campylobacter spec	Yes				822
Cancer deaths, total (per 100,000)	Overall cancer death rate	Yes	Yes	Yes		486
Cancer prevalence: adults (percent)	Respondent-reported prevalence of cancer among adults aged 18 and over	Yes				50045
Cancer survival (percent)	Percent of cancer survivors who are living 5 years or longer after diagnosis	Yes				498
Carbon monoxide poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that ca	Yes				719
Cardiac surgery patients with controlled 6 AM postoperative blood glucose (percent)	Hospital Compare: Cardiac surgery patients with controlled 6 AM postoperative blood glucose	Yes	Yes	Yes	Yes	339
Cardiovascular death: hemodialysis patients (per 1,000 patients years at risk)	Cardiovascular death rate for persons on hemodialysis	Yes				465
Cardiovascular death: kidney transplant patients (per 1,000 patients years at risk)	Cardiovascular death rate for persons with a functioning kidney transplant	Yes				467
Catheter use for vascular access: hemodialysis patients (percent)	Percent of adult hemodialysis patients who use catheters as the only mode of vascular access	Yes				470
Central line-associated bloodstream infections (Standardized Infection Ratio (SIR))	Central line-associated bloodstream infections	Yes				851
Certificate programs in public health: 2-year colleges (percent)	Percent of 2-year colleges that offer public health or related Certificate programs	Yes				1389
Certificates of birth, use of latest standard form: states (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States	Yes				1382
Certificates of death, use of latest standard form: states (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States	Yes				1383
Cervical cancer screening: women 21-65 years (percent)	Percent of women who receive a cervical cancer screening based on the most recent guidelines	Yes				505
Cesarean delivery, first (percent)	Percent of live births that are cesarean deliveries among low-risk (full-term, singleton, vertex presenta	Yes				1133
Cesarean delivery, repeat (percent)	Percent of live births that are cesarean among low-risk (full-term, singleton, vertex presentation) won	Yes				1134
Child developmental concerns addressed by doctors: parents (percent)	Percent of parents who receive information from their doctors or other Health Care professionals whe	Yes				741
Child maltreatment deaths (per 100,000)	Child maltreatment death rate	Yes				1085
Child maltreatment, nonfatal (per 1,000)	Nonfatal child maltreatment	Yes				1084
Children's Hospitals (number)	Number of Children's Hospitals participating in the Medicare program	Yes	Yes		Yes	10034
Chlamydia (per 100,000)	Chlamydia rate	Yes	Yes	Yes		20
Chlamydia cases (number)	Number of reported cases of chlamydia		Yes	Yes		10064
Chlamydia screening: females 16-20 years with Medicaid (percent)	Percent of sexually active females aged 16 to 20 years enrolled in Medicaid plans who are screened fo	Yes				1486
Chlamydia screening: females 16-20 years with private health insurance (percent)	Percent of sexually active females aged 16 to 20 years enrolled in commercial health insurance plans w	Yes				1488
Chlamydia screening: females 21-24 years with Medicaid (percent)	Percent of sexually active females aged 21 to 24 years enrolled in Medicaid plans who are screened fo	Yes				1487
Chlamydia screening: females 21-24 years with private health insurance (percent)	Percent of sexually active females aged 21 to 24 years enrolled in commercial health insurance plans w	Yes				1489
Chlamydia: females <24 years in a national job training program (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among females aged ;	Yes				1476
Chlamydia: females 15-24 years attending family planning clinics (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among females aged :	Yes				1475
Chlamydia: males <24 years in a national job training program (percent)	Percent of adolescents and young adults with Chlamydia trachomatis infections-Among males aged 24	Yes				1477
Chlordane level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				700
Chlorpyrifos level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				704
Cholesterol levels: adults (mean)	Mean total blood cholesterol levels among adults	Yes				884
Cholesterol management (percent)	Managed low-density lipoprotein cholesterol (LDL-C) among hyperlipidemic adults aged =20 years.	Yes				50073
Cholesterol screening: adults (percent)	Percent of adults who have had their blood cholesterol checked within the preceding 5 years	Yes				886
Chronic diseases prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services,	Yes				602
Chronic hepatitis B: children 1-24 months (number)	Number of chronic hepatitis B virus infections in infants and young children (perinatal infections)	Yes				962
Chronic kidney disease deaths: adults (per 100)	Death rate among adults 18 years or older with chronic kidney disease	Yes				484
Chronic kidney disease Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have chronic kidney disease	Yes	Yes	Yes	Yes	287
Chronic kidney disease Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have chronic kidney disease	Yes	Yes	Yes	Yes	288
Chronic kidney disease: adults (percent)	Percent of the adults with chronic kidney disease	Yes				483
Chronic lower respiratory disease (CLRD) deaths (per 100,000)	Chronic lower respiratory disease (CLRD) death rate	Yes	Yes	Yes		50012
Chronic obstructive pulmonary disease (COPD) Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have chronic obstructive pulmonary disease	Yes	Yes	Yes	Yes	289
Chronic obstructive pulmonary disease (COPD) Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have chronic obstructive pulmonary disease	Yes	Yes	Yes	Yes	290
Chronic obstructive pulmonary disease admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Chronic obstructive pulmonary disease admission rate among Medicare	Yes	Yes	Yes	Yes	2039
Cigar smoking: adults (percent)	Percent of adults aged 18 years and over smoking cigars	Yes				1500
Cigar use in past month: teens (percent)	Percent of adolescent reporting use of cigars in the past month - Students (grades 9 through 12)	Yes	Yes			1504
Cigar use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating cigar use in the past 12 months	Yes				1512

Cigar use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating cigar use in past 12 months	Yes				1508
Cigarette abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Cigarette smoking	Yes				1144
Cigarette smoking: adults (percent)	Percent of adults aged 18 years and over smoking cigarettes	Yes				1498
Cigarette smoking: adults (percent) (Source: BRFSS)	Percent of adults 18 years and over that report currently smoking cigarettes		Yes	Yes		13
Cigarette use in past month: teens (percent)	Percent of adolescent reporting use of cigarettes in the past month - Students (grades 9 through 12)	Yes	Yes			1502
Cigarette use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating cigarette use in the past 12 months	Yes				1510
Cigarette use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating cigarette use in past 12 months	Yes				1506
Cirrhosis deaths (per 100,000)	Cirrhosis death rate	Yes	Yes			1425
Clean hands, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Clean: Wash hands and surfaces often	Yes				835
Clinical Laboratory Improvement Act (CLIA) Providers (number)	Number of Clinical Laboratory Improvement Act (CLIA) providers participating in the Medicare program	Yes	Yes		Yes	10035
Clinical preventive measure use: females 65+ years (percent)	Percent of older adults who are up to date on a core set of clinical preventive measures-Females	Yes				1262
Clinical preventive measure use: males 65+ years (percent)	Percent of older adults who are up to date on a core set of clinical preventive measures-Males	Yes				1261
Cochlear implant surgeries: deaf persons (per 10,000)	Rate of persons who are deaf or very hard of hearing and who have new cochlear implants	Yes				745
College degree: adults 25+ (percent)	Percent of population age 25+ with 4-year college degree or higher	Yes	Yes	Yes		34
Colonoscopy use: adults 50-75 (percent) (Source: NHIS)	Percent of adults aged 50-75 who have had a colonoscopy in the past 10 years	Yes				50064
Colonoscopy/sigmoidoscopy: adults 50+ (percent) (Source: BRFSS)	Percent of adults age 50+ that report ever having had a sigmoidoscopy or colonoscopy		Yes	Yes		117
Colorectal cancer deaths (per 100,000)	Colorectal cancer death rate per 100,000	Yes	Yes	Yes		86
Colorectal cancer deaths, including unspecified sites (per 100,000)	Colorectal cancer death rate, including unspecified sites, per 100,000	Yes				490
Colorectal cancer Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have colorectal cancer	Yes	Yes	Yes	Yes	301
Colorectal cancer Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have colorectal cancer	Yes	Yes	Yes	Yes	302
Colorectal cancer screening: persons 50-75 years (percent)	Percent of adults who receive a colorectal cancer screening based on the most recent guidelines	Yes				506
Commitment from responsible or responding party to reduce risk: news stories on foodborne disease (percent)	Percent of crisis and emergency risk messages embedded in print and broadcast news stories that express commitment	Yes				10099
Community Mental Health Centers (number)	Number of Community Mental Health Centers participating in the Medicare program	Yes	Yes		Yes	10036
Competencies for public health professionals: Local agencies (percent)	Percent of Federal, Tribal, State, and local public health agencies that incorporate core competencies for public health professionals	Yes				1372
Complete vaccination: children 19-35 months (percent)	Percent of children aged 19 to 35 months who receive the recommended doses of DTaP, polio, MMR, Hib, and Hepatitis B	Yes	Yes			1008
Complex activity limitation: adults (percent)	Percent of adults aged 18 and over with at least one complex activity limitation	Yes				50043
Comprehensive care systems: children with special needs (percent)	Percent of children with special health care needs who receive their care in family-centered, comprehensive care systems	Yes				1154
Comprehensive care systems: teens with special needs (percent)	Percent of children with special health care needs who receive their care in family-centered, comprehensive care systems	Yes				1155
Comprehensive Outpatient Rehabilitation Facilities (number)	Number of Comprehensive Outpatient Rehabilitation (CORF) Facilities participating in the Medicare program	Yes	Yes		Yes	10037
Condition not improving: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems in the past 12 months who reported their condition not improving	Yes				766
Condom & hormonal/intrauterine contraception use at first intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse	Yes				798
Condom & hormonal/intrauterine contraception use at first intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at first intercourse	Yes				799
Condom & hormonal/intrauterine contraception use at last intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse	Yes				800
Condom & hormonal/intrauterine contraception use at last intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom and hormonal or intrauterine contraception at last intercourse	Yes				801
Condom use at first intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom at first intercourse	Yes				794
Condom use at first intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom at first intercourse	Yes				795
Condom use at last intercourse: females 15-19 years (percent)	Percent of sexually active females aged 15 to 19 years who use a condom at last intercourse	Yes				796
Condom use at last intercourse: males 15-19 years (percent)	Percent of sexually active males aged 15 to 19 years who use a condom at last intercourse	Yes				797
Condom use: unmarried females 15-44 years (percent)	Percent of sexually active persons who use condoms-Unmarried females aged 15 to 44 years	Yes				940
Condom use: unmarried males 15-44 years (percent)	Percent of sexually active persons who use condoms-Unmarried males aged 15 to 44 years	Yes				941
Congenital Rubella Syndrome (number)	Number of Congenital Rubella Syndrome (CRS) among children aged 1 year and under (U.S.-acquired)	Yes				978
Congestive heart failure admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Congestive heart failure admission rate among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2041
Congregate care: adults with disabilities (number)	Number of adults with disabilities (aged 22 years and older) living in congregate care facilities with 16 or more residents	Yes				524
Congregate care: persons <22 years with disabilities (number)	Number of children and youth with disabilities (aged 21 years and under) living in congregate care facilities with 16 or more residents	Yes				525
Consultation on work related asthma (percent)	Percent of adults with current asthma who have discussed with a doctor or other health professional about work-related asthma	Yes				1420
Contraceptive services/supplies need and receipt through public sector (percent)	Percent of females in need of publicly supported contraceptive services and supplies who receive those services and supplies	Yes				819
Contraceptive use at most recent sexual intercourse (percent)	Percent of females or their partners at risk of unintended pregnancy who used contraception at most recent sexual intercourse	Yes				785
Cook to proper temperature, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Cook: Cook to proper temperatures	Yes				837

COPD deaths: adults 45+ years (per 100,000)	Chronic obstructive pulmonary disease (COPD) death rate among adults aged 45 years and over	Yes				1406
COPD ED visits: adults 45+ years (per 10,000)	Rate of hospital emergency department visits for chronic obstructive pulmonary disease (COPD) among	Yes				1423
COPD hospitalizations: adults 45+ years (per 10,000)	Hospitalization rate for chronic obstructive pulmonary disease (COPD) among adults aged 45 years and	Yes				1422
Core Competencies in Public Health in curricula (percent)	Percent of Council on Education for Public Health (CEPH) accredited schools of public health, CEPH acc	Yes				1373
Coronary heart disease deaths (per 100,000)	Coronary heart disease death rate	Yes	Yes	Yes		877
Critical Access Hospitals (number)	Number of Critical Access Hospitals participating in the Medicare program	Yes	Yes		Yes	10038
D.O. training, counseling for health promotion (percent)	Percent of DO-granting medical schools that include training in counseling for health promotion and di	Yes				612
D.O. training, cultural diversity (percent)	Percent of DO-granting medical schools that include training in cultural diversity	Yes				613
D.O. training, environmental health (percent)	Percent of DO-granting medical schools that include training in environmental health	Yes				615
D.O. training, evaluation of health sciences literature (percent)	Percent of DO-granting medical schools that include training in evaluation of health sciences literatur	Yes				614
D.O. training, global health (percent)	Percent of DO-granting medical schools that include training in global health in health	Yes				617
D.O. training, public health systems (percent)	Percent of DO-granting medical schools that include training in public health systems	Yes				616
DDT level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				701
Deaf or a lot of trouble hearing: adults (percent)	Percent of adults aged 18 and over who have a lot of trouble hearing or who are deaf	Yes				50062
Death: kidney transplant patients (per 1,000 patients years at risk)	Total death rate for persons with a functioning kidney transplant	Yes				466
Deaths, all causes (per 100,000)	Death rate per 100,000 population from all causes of death	Yes	Yes	Yes		71
Deaths: adults 20-24 years (per 100,000)	Rates of adolescent and young adult deaths-Young adults aged 20 to 24 years	Yes	Yes			1128
Deaths: children 10-14 years (per 100,000)	Rates of adolescent and young adult deaths-Adolescents aged 10 to 14 years	Yes	Yes			1126
Deaths: children 1-4 years (per 100,000)	Rate of child deaths-Children aged 1 to 4 years	Yes	Yes			1124
Deaths: children 5-9 years (per 100,000)	Rate of child deaths-Children aged 5 to 9 years	Yes	Yes			1125
Deaths: infants with Down syndrome (per 1,000 infants with Down syndrome)	One-year mortality rate for infants with Down syndrome	Yes				1187
Deaths: teens 15-19 years (per 100,000)	Rates of adolescent and young adult deaths-Adolescents aged 15 to 19 years	Yes	Yes			1127
Dehydration admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Dehydration admission rate among Medicare fee-for-service beneficiari	Yes	Yes	Yes	Yes	2042
Dental care not received due to cost (percent)	Percent of persons of all ages who did not get dental care in the previous 12 months due to cost	Yes	Yes			50067
Dental care programs: school-based health centers (percent)	Percent of school-based health centers with an oral health component that includes dental care	Yes				1285
Dental caries experience: children 3-5 years (percent)	Percent of young children aged 3 to 5 years with dental caries experience in their primary teeth	Yes				1269
Dental caries experience: children 6-9 years (percent)	Percent of children aged 6 to 9 years with dental caries experience in their primary and permanent tee	Yes				1270
Dental caries experience: teens 13-15 years (percent)	Percent of adolescents aged 13 to 15 years with dental caries experience in their permanent teeth	Yes				1271
Dental decay, untreated: adults 35-44 years (percent)	Percent of adults aged 35 to 44 years with untreated dental decay	Yes				1275
Dental decay, untreated: adults 65-74 years (percent)	Percent of older adults aged 65 to 74 years with untreated coronal caries	Yes				1276
Dental decay, untreated: adults 75+ years (percent)	Percent of older adults aged 75 years and older with untreated root surface caries	Yes				1277
Dental decay, untreated: children 3-5 years (percent)	Percent of young children aged 3 to 5 years with untreated dental decay in their primary teeth	Yes				1272
Dental decay, untreated: children 6-9 years (percent)	Percent of children aged 6 to 9 years with untreated dental decay in their primary and permanent teet	Yes				1273
Dental decay, untreated: teens 13-15 years (percent)	Percent of adolescents aged 13 to 15 years with untreated dental decay in their permanent teeth	Yes				1274
Dental examinations, annual: adults with diabetes (percent)	Percent of persons with diagnosed diabetes who have at least an annual dental examination	Yes				553
Dental professional directing dental program: IHS programs (number)	Number of Indian Health Service Areas and Tribal health programs that serve jurisdictions of 30,000 or	Yes				1292
Dental professional directing dental program: state & local agencies (percent)	Percent of States (including the District of Columbia) and local health agencies that serve jurisdictions	Yes				1291
Dental sealant programs: school-based health centers (percent)	Percent of school-based health centers with an oral health component that includes dental sealants	Yes				1284
Dental sealants: children 3-5 years (percent)	Percent of children aged 3 to 5 years who have received dental sealants on one or more of their prim	Yes				1281
Dental sealants: children 6-9 years (percent)	Percent of children aged 6 to 9 years who have received dental sealants on one or more of their perm	Yes				1282
Dental sealants: teens 13-15 years (percent)	Percent of adolescents aged 13 to 15 years who have received dental sealants on one or more of their	Yes				1283
Dental service use: low-income children (percent)	Percent of low-income children and adolescents who received any preventive dental service during th	Yes				1267
Dental visits: ages 2+ (percent) (Source: NHIS)	Percent of persons 2 years of age and over who had have a dental visit in the past year	Yes				50053
Dentist rate (per 100,000)	The total number of active dentists per capita		Yes	Yes		130
Dentist training, counseling for health promotion (percent)	Percent of Doctor of Dental Surgery and/or Doctor of Dental Medicine granting colleges and schools of	Yes				10081
Dentist training, cultural diversity (percent)	Percent of Doctor of Dental Surgery and/or Doctor of Dental Medicine granting colleges and schools of	Yes				10082
Dentist training, environmental health (percent)	Percent of Doctor of Dental Surgery and/or Doctor of Dental Medicine granting colleges and schools of	Yes				10084
Dentist training, evaluation of health sciences literature (percent)	Percent of Doctor of Dental Surgery and/or Doctor of Dental Medicine granting colleges and schools of	Yes				10083

Dentist training, global health (percent)	Percent of Doctor of Dental Surgery and/or Doctor of Dental Medicine granting colleges and schools of					10086
Dentist training, public health systems (percent)	Percent of Doctor of Dental Surgery and/or Doctor of Dental Medicine granting colleges and schools of					10085
Depression Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have depression	Yes	Yes	Yes	Yes	291
Depression Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have depression	Yes	Yes	Yes	Yes	292
Depression screening by primary care providers: adults (percent)	Percent of primary care physician office visits that screen adults aged 19 years and older for depression	Yes				1120
Depression screening by primary care providers: teens (percent)	Percent of primary care physician office visits that screen youth aged 12 to 18 years for depression	Yes				1121
Diabetes deaths (per 100,000)	Diabetes mellitus death rate per 100,000	Yes	Yes	Yes		50011
Diabetes long-term complications admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Diabetes long-term complications admission rate among Medicare fee-f	Yes	Yes	Yes	Yes	2038
Diabetes management benefit use: older adults with diabetes (percent)	Percent of older adults who receive Diabetes Self- Management Benefits	Yes				1263
Diabetes Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have diabetes	Yes	Yes	Yes	Yes	293
Diabetes Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have diabetes	Yes	Yes	Yes	Yes	294
Diabetes, new cases (per 1,000)	Rate of newly diagnosed diabetes among adults aged 18-84	Yes				555
Diabetes: adults (percent)	Percent of adults that report having been diagnosed as having diabetes		Yes	Yes		125
Diabetes-related deaths (per 100,000)	Diabetes death rate	Yes				547
Diagnosed diabetes: adults with diabetes (percent)	Percent of adults aged 20 years and older with diabetes whose condition has been diagnosed	Yes				546
Dialysis Medicare users (count)	Medicare fee-for-service beneficiaries who used dialysis services	Yes	Yes	Yes	Yes	50005
Dialysis Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used dialysis services	Yes	Yes	Yes	Yes	50006
Dialysis Medicare visits (per 1,000 beneficiaries)	Rate of dialysis visits among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	50004
Dilated eye examinations: adults (percent)	Percent of adults who have a comprehensive eye examination, including dilation, within the past 2 yea	Yes				1566
Dioxin level: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				708
Disapproval of heavy drinking: 10th graders (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-10th gra	Yes				1445
Disapproval of heavy drinking: 8th graders (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-8th gra	Yes				1444
Disapproval of heavy drinking: high school seniors (percent)	Percent of adolescents who disapprove of having one or two alcoholic drinks nearly every day-12th gra	Yes				1446
Disapproval of trying marijuana: 10th graders (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-10th graders	Yes				1448
Disapproval of trying marijuana: 8th graders (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-8th graders	Yes				1447
Disapproval of trying marijuana: high school seniors (percent)	Percent of adolescents who disapprove of trying marijuana or hashish once or twice-12th graders	Yes				1449
Disorders diagnosed through newborn blood spot screening (percent)	Percent of children diagnosed with a disorder through newborn blood spot screening who experience	Yes				1167
Domestic water use (gallons)	Per capita domestic water withdrawals with respect to use and conservation	Yes				673
Drownings (per 100,000)	Drowning death rate	Yes	Yes	Yes		1081
Drug poisoning deaths (per 100,000)	Rate per 100,000 of deaths with underlying causes of drug related poisonings	Yes	Yes	Yes		10016
Drug-induced deaths (per 100,000)	Drug-induced death rate	Yes	Yes			1426
DTaP vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-4 doses diphtheria-te	Yes	Yes			993
DTaP vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-DTaP vaccine (4 or more doses)	Yes				1003
Durable medical equipment Medicare service events (per 1,000 beneficiaries)	Rate of durable medical equipment service events among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2026
Durable medical equipment Medicare users (count)	Medicare fee-for-service beneficiaries who used durable medical equipment services	Yes	Yes	Yes	Yes	257
Durable medical equipment Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used durable medical equipment services	Yes	Yes	Yes	Yes	258
Ear infections, 3+ in 12 months: children (percent)	Percent of children under age 18 years who have had three or more ear infections in the last 12 mont	Yes				50038
Early death: hemodialysis patients (per 1,000 patients years at risk)	Death rate in hemodialysis patients within the first 3 months of initiation of renal replacement therap	Yes				464
Early infant diagnostic hearing evaluation (percent)	Percent of infants who did not pass the hearing screening that received audiologic evaluation at no lat	Yes				757
Early intervention services: children with disabilities (percent)	Percent of children with disabilities, birth through age 2 years, who receive early intervention services	Yes				539
Early intervention services: infants with hearing loss (percent)	Percent of infants with confirmed hearing loss enrolled for intervention services no later than age 6 m	Yes				758
Eating disorders: teens (percent)	Percent of adolescents who engage in disordered eating behaviors in an attempt to control their weigl	Yes	Yes			1108
E-code collection at hospital discharge: states (percent)	Percent of states and the District of Columbia with statewide hospital discharge data systems that rou	Yes				1094
E-code collection in emergency department: states (percent)	Percent of states and the District of Columbia with statewide emergency department data systems tha	Yes				1093
Elder abuse, neglect, and exploitation information: States (number)	Number of States-and the District of Columbia-that collect and make publicly available information on	Yes				1256
Electronic health records in medical practices (percent)	Percent of medical practices that use electronic health records	Yes				872
Emergency department visit rate (per 1,000 beneficiaries)	Rate among beneficiaries of inpatient or hospital outpatient emergency department visits	Yes	Yes	Yes	Yes	282
Emergency department visits (count)	Total inpatient or hospital outpatient emergency department visits	Yes	Yes	Yes	Yes	281

Emergency personnel response time (minutes)	The time necessary to activate designated personnel in response to public health emergency in minute	Yes				1394
Empathy about health risk expressed: news stories on foodborne disease outbreaks or national disaster	Percent of crisis and emergency risk messages embedded in print and broadcast news stories that express	Yes				10098
Employment at indoor workplaces that prohibit smoking (percent)	Percent of persons covered by indoor worksite policies that prohibit smoking	Yes				1492
Employment: persons 16-64 years with disabilities (percent)	Employment rates among people with disabilities	Yes				516
Employment: persons with serious mental illness (percent)	Percent of persons with serious mental illness (SMI) who are employed	Yes				1116
End Stage Renal Disease Facilities (number)	Number of End Stage Renal Disease (ESRD) facilities participating in the Medicare program	Yes	Yes		Yes	10039
End-stage kidney failure due to diabetes (per 1,000,000)	Incidence rate of kidney failure due to diabetes	Yes				475
End-stage kidney failure: persons with diabetes (per 1,000,000)	Incidence rate of kidney failure due to diabetes among persons with diabetes	Yes				476
End-stage renal disease (per 1,000,000)	Incidence rate of end-stage renal disease (ESRD)	Yes				462
End-stage renal disease deaths: hemodialysis patients (per 1,000 patients years at risk)	Total death rate for persons on hemodialysis	Yes				463
Enforcement of illegal tobacco sales to minors laws: States (number)	Illegal tobacco sales rate of <=5% to minors through enforcement of laws prohibiting the sale of tobacco	Yes				1493
Enforcement of illegal tobacco sales to minors laws: Territories (number)	Illegal tobacco sales rate of <=5% to minors through enforcement of laws prohibiting the sale of tobacco	Yes				1494
Enrollment in college immediately after high school: high school completers (percent)	Percent of high school completers who were enrolled in college the October immediately after completion	Yes				10071
Epidemiology services: Local public health agencies (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology	Yes				1363
Epidemiology services: State public health agencies (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidemiology	Yes				1362
Escherichia coli infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction of Shiga toxin	Yes				823
Estimated Poverty (percent) (Source: SAIPE)	Small area estimates of population living below the poverty line	Yes	Yes	Yes		10004
Excessive drinking: adults (percent)	Percent of adults who drank excessively in the previous 30 days	Yes				1458
Excessive drinking: adults (percent) (Source: BRFSS)	Percent of adults that report excessive drinking, either chronic high alcohol consumption or binge drinking		Yes	Yes		150
Extracurricular activities: teens aged 12-17 (percent)	Percent of adolescents who participate in extracurricular and out-of-school activities	Yes				404
Eye examinations, annual: adults with diabetes (percent)	Percent of adults with diabetes who have an annual dilated eye examination	Yes				551
Fair or poor health (percent) (Source: NHIS)	Percent of persons of all ages with self-, or proxy-, reported fair or poor health	Yes				50048
Fair or poor health: adults (percent) (Source: BRFSS)	Percent of adults that report fair or poor health		Yes	Yes		5
Fall deaths (per 100,000)	Fall-related death rate: All persons	Yes	Yes	Yes		1079
Fall deaths: older adults (per 100,000)	Fall-related death rate: Adults aged 65 years and over	Yes				1080
Fall ED visits: older adults (per 100,000)	Rate of emergency department visits due to falls among older adults	Yes				1259
Falling: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems who have fallen in the past 5 years while experiencing	Yes				769
Family planning clinics offering all contraception methods (percent)	Percent of publicly funded family planning clinics that offer the full range of FDA-approved methods of family planning	Yes				781
Family planning clinics offering emergency contraception (percent)	Percent of publicly funded family planning clinics that offer emergency contraception onsite	Yes				782
Family planning services income eligibility: states (number)	Number of States that set the income eligibility level for Medicaid-covered family planning services to <=150%	Yes				818
Federally Qualified Health Center and Rural Health Clinic Medicare users (count)	Medicare fee-for-service beneficiaries who used Federally Qualified Health Center and Rural Health Clinic	Yes	Yes	Yes	Yes	50002
Federally Qualified Health Center and Rural Health Clinic Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used Federally Qualified Health Center and Rural Health Clinic	Yes	Yes	Yes	Yes	50003
Federally Qualified Health Center and Rural Health Clinic visits (per 1,000 beneficiaries)	Rate of Federally Qualified Health Center and Rural Health Clinic visits among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	50001
Federally Qualified Health Centers (number)	Number of Federally Qualified Health Centers (FQHC) participating in the Medicare program	Yes	Yes		Yes	10040
Female Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who are female	Yes	Yes	Yes	Yes	309
Fetal alcohol syndrome (per 10,000 live births)	Rate of fetal alcohol syndrome (FAS)	Yes				1146
Fetal deaths (per 1,000 live births plus fetal deaths)	Rate of fetal deaths at 20 or more weeks of gestation	Yes				1156
Few fruits/vegetables: adults (percent)	Adults that report fewer than 5 servings of fruits/vegetables per day		Yes	Yes		121
Fibrinolytic therapy timeliness: heart attack patients (percent)	Percent of eligible patients with heart attacks who receive fibrinolytic therapy within 30 minutes of hospital arrival	Yes				878
Firearm-related deaths (per 100,000)	Firearm-related death rate	Yes	Yes	Yes		1042
Flu vaccination: adults 65+ (percent) (Source: BRFSS)	Percent of adults age 65+ that report having Influenza vaccine in past 12 months		Yes	Yes		119
Fluoridation of community water (percent)	Percent of the U.S. population served by community water systems with optimally fluoridated water	Yes	Yes			1265
Fluoride programs: school-based health centers (percent)	Percent of school-based health centers with an oral health component that includes topical fluoride	Yes				1286
Folic acid intake: reproductive-aged women (percent)	Percent of women of childbearing potential with intake of at least 400 µg of folic acid from fortified foods	Yes				1171
Follow-up to newborn screening (percent)	Percent of screen-positive children who receive follow-up testing within the recommended time period	Yes				1174
Food allergy: children (percent)	Percent of children under age 18 years who have food allergies	Yes				50035
Food insecurity: households (percent)	Percent of households with food insecurity	Yes				1243
Food insecurity: households with children (percent)	Percent of household with children with very low food security	Yes				1228

Foot examinations, annual: adults with diabetes (percent)	Percent of adults with diabetes who have at least an annual foot examination	Yes	Yes	552
Formula supplementation: breastfed newborns (percent)	Percent of breastfed newborns who receive formula supplementation within the first 2 days of life	Yes		1185
Free school lunch eligibility (percent)	Percent of total students eligible to participate in the National School Lunch Program under the Free L		Yes Yes	10005
Fruit consumption (cup equivalents per 1,000 calories)	Mean daily intake of fruit (in cup equivalents per 1,000 calories) by persons aged 2 years and over	Yes		1217
Fruit/vegetable availability required: school districts (percent)	Percent of school districts that require schools to make fruits or vegetables available whenever other f	Yes		10066
Functional limitations: older adults (percent)	Percent of older adults 65 years and over who have moderate to severe functional limitations	Yes		1245
Genetic counseling: women with family history of breast/ovarian cancer (percent)	Percent of women with a family history of breast and/or ovarian cancer who receive genetic counselin	Yes		850
Genital herpes simplex type 2: adults 20-29 years (percent)	Percent of young adults with genital herpes infection due to herpes simplex type 2	Yes		1482
Geriatric dentists (percent)	Percent of the health care workforce with geriatric certification-Dentists	Yes		1253
Geriatric dietitians (percent)	Percent of the health care workforce with geriatric certification-Registered dietitians	Yes		1255
Geriatric nurses (percent)	Percent of the health care workforce with geriatric certification-Registered nurses	Yes		1252
Geriatric physical therapists (percent)	Percent of the health care workforce with geriatric certification-Physical therapists	Yes		1254
Geriatric physicians (percent)	Percent of the health care workforce with geriatric certification-Physicians	Yes		1250
Geriatric psychiatrists (percent)	Percent of the health care workforce with geriatric certification-Geriatric psychiatrists	Yes		1251
Giardiasis (number)	Number of reported cases of giardiasis	Yes	Yes Yes	10021
Gini coefficient (ratio)	Gini coefficient of household income inequality	Yes	Yes Yes	37
Global Disease Detection increased testing capacity (number)	Diagnostic testing capacity in host countries and regionally through the Global Disease Detection (GDD)	Yes		848
Global Disease Detection Regional Centers (number)	Number of Global Disease Detection (GDD) Regional Centers worldwide to detect and contain emergir	Yes		846
Global Disease Detection training of public health professionals (number)	Number of public health professionals trained by Global Disease Detection: Global Disease Detection (Yes		847
Glucose A1c < 7%: adults with diabetes (percent)	Percent of adults with diagnosed diabetes with an HbA1c value less than 7 percent	Yes		559
Glucose A1c > 9%: adults with diabetes (percent)	Percent of adults with diagnosed diabetes with an HbA1c value greater than 9 percent	Yes		558
Glucose A1c test, biannual: adults with diabetes (percent)	Percent of adults with diabetes who have a glycosylated hemoglobin measurement at least twice a ye	Yes	Yes	550
Glucose monitoring, daily: adults with diabetes (percent)	Percent of adults with diabetes who perform self-blood glucose-monitoring at least once daily	Yes	Yes	554
Gonorrhea rate (per 100,000)	Rate of gonorrhea cases per 100,000 population	Yes	Yes Yes	10062
Gonorrhea: females 15-44 years (per 100,000)	Gonorrhea rate among females aged 15 to 44 years	Yes		1478
Gonorrhea: males 15-44 years (per 100,000)	Gonorrhea rate among males aged 15 to 44 years	Yes		1479
Graduated driver licensing laws: states (number)	Number of states and the District of Columbia with "good" graduated driver licensing (GDL) laws	Yes		1058
Group B streptococcal disease: newborns (per 1,000)	Early onset group B streptococcal disease rate	Yes		973
Hay fever/respiratory allergy: children (percent)	Percent of children under age 18 who have hay fever or other respiratory allergies	Yes		50037
Hazardous material plan: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes		685
Hazardous sites posing risk (number)	Number of hazardous sties that pose risks to human health and the environment	Yes		677
HbA1c test: diabetic Medicare beneficiaries 65-75 years (percent)	Percent of diabetic Medicare enrollees ages 65-75 receiving a hemoglobin A1c test in the past year		Yes Yes	29
Headache or migraine, severe: adults (percent)	Percent of adults aged 18 and over with pain during the past 3 months due to severe headache or mig	Yes		50039
Health care evaluation: adults with balance/dizziness problems (percent)	Percent of adults with balance or dizziness problems in the past 12 months who have ever seen a Heal	Yes		762
Health care evaluation: adults with tinnitus (percent)	Percent of adults bothered by tinnitus in the past 12 months who have seen a doctor	Yes		759
Health care organizations monitoring adverse events (percent)	Percent of health care organizations that are monitoring and analyzing adverse events associated with	Yes		1196
Health care provider explaining things (percent)	Percent of persons who report that their Health Care provider always explained things so they could u	Yes		854
Health care provider listening skills (percent)	Percent of persons who report that their Health Care provider always listened carefully to them	Yes		853
Health care provider referrals to hearing specialists (per 1,000)	Rate of persons who are referred by their primary care physician or other health care provider for hea	Yes		751
Health care provider showing respect (percent)	Percent of persons who report that their Health Care provider always showed respect for what they h	Yes		855
Health care provider spending enough time with patient (percent)	Percent of persons who report that their health care provider always spent enough time with them	Yes		856
Health ed. certification requirements: elementary school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, Yes			731
Health ed. certification requirements: high school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, Yes			733
Health ed. certification requirements: middle school (percent)	Percent of schools that require newly hired staff who teach required health instruction to be certified, Yes			732
Health ed. goals, advocacy: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes		638
Health ed. goals, communication skills: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes		642
Health ed. goals, concept comprehension: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes		636
Health ed. goals, decision-making skills: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes		641

Health ed. goals, external influences: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes			639
Health ed. goals, healthy behaviors: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes			640
Health ed. goals, information access: schools (percent)	Percent of elementary, middle, and senior high schools that have health education goals or objectives	Yes			637
Health ed. teachers trained in behavior change (percent)	Percent of required health education classes or courses with a teacher who has had professional devel	Yes			737
Health ed. training requirements: elementary schools (percent)	Percent of schools that require newly hired staff who teach required health education to have undergr	Yes			728
Health ed. training requirements: high schools (percent)	Percent of schools that require newly hired staff who teach required health education to have undergr	Yes			730
Health ed. training requirements: middle schools (percent)	Percent of schools that require newly hired staff who teach required health education to have undergr	Yes			729
Health education, alcohol/drug use: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			571
Health education, all priority areas: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			566
Health education, benefits of sleep: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			656
Health education, growth & development: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			654
Health education, hand washing info: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			652
Health education, health screening: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			658
Health education, injury prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			567
Health education, nutrition: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			573
Health education, oral health: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			653
Health education, physical activity: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			574
Health education, pregnancy, HIV/AIDS, STD: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			572
Health education, suicide prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			569
Health education, tobacco use: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			570
Health education, violence prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide comprehensive school health edu	Yes			568
Health education, vision & hearing loss prevention: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			657
Health educations, sun safety: schools (percent)	Percent of elementary, middle, and senior high schools that provide school health education to promo	Yes			655
Health information, alcohol/drug: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on alcohol ar	Yes			580
Health information, HIV & STD: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on HIV, AIDS	Yes			582
Health information, injury: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on unintentic	Yes			576
Health information, nutrition: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on unhealthy	Yes			583
Health information, physical activity: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on inadequal	Yes			584
Health information, suicide: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on suicide	Yes			578
Health information, tobacco use: post-secondary schools (percent)	percent of college and university students who receive information from their institution on tobacco u	Yes			579
Health information, unintended pregnancy: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on unintend	Yes			581
Health information, violence: post-secondary schools (percent)	Percent of college and university students who receive information from their institution on violence	Yes			577
Health instruction meets time standards: elementary schools (percent)	Percent of schools that require cumulative instruction in health education that meet the US National F	Yes			734
Health instruction meets time standards: high schools (percent)	Percent of schools that require cumulative instruction in health education that meet the US National F	Yes			736
Health instruction meets time standards: middle schools (percent)	Percent of schools that require cumulative instruction in health education that meet the US National F	Yes			735
Health insurance coverage, private through workplace: persons <65 (percent)	Percent of persons under age 65 who have private health insurance coverage obtained through the wc	Yes			50056
Health insurance, none: <65 years (percent) (Source: SAHIE)	Percent of persons under 65 years of age without health insurance		Yes	Yes	23
Health insurance, none: persons <65 (percent) (Source: NHIS)	Percent of persons under age 65 with no health insurance coverage	Yes			50058
Health insurance: <65 years (percent)	Percent of persons under age 65 years with health (medical) insurance	Yes			368
Health promotion programs for caregivers: States (number)	States and the District of Columbia with health promotion programs for caregivers	Yes			529
Health promotion programs for people with disabilities: States (number)	States and the District of Columbia with health promotion programs for people with disabilities	Yes			527
Health provider seen: adults with joint-related symptoms (percent)	Percent of adults with chronic joint symptoms who have seen a health care provider for their symptom	Yes			425
Health websites meeting 3+ criteria to assess reliability (percent)	Percent of health-related Websites that meet three or more evaluation criteria disclosing information	Yes			863
Healthy food outlets (percent)	Percent of zip codes in county with a healthy food outlet			Yes	48
Healthy weight prior to pregnancy (percent)	Percent of women delivering a live birth who had a healthy weight prior to pregnancy	Yes			1182
Healthy weight: adults (percent)	Percent of adults who are at a healthy weight	Yes			1207
Hearing aid use: adults 20-69 years (per 1,000)	Rate of adults aged 20 to 69 years with hearing loss who have ever used a hearing aid	Yes			744
Hearing loss, noise-induced, work-related (per 10,000 full-time workers)	Rate of work-related, noise-induced hearing loss	Yes			1302

Hearing loss, noise-induced: adults (per 1,000)	Rate of adults who have elevated hearing thresholds, or audiometric notches, in high frequencies (3, 4 Yes	755
Hearing loss, noise-induced: teens (per 1,000)	Rate of adolescents who have elevated hearing thresholds, or audiometric notches, in high frequency: Yes	754
Heart attack awareness: importance of calling 9-1-1 (percent)	Percent of adults aged 20 years and over who are aware of the importance of accessing rapid emerger Yes	896
Heart attack awareness: symptoms (percent)	Percent of adults aged 20 years and over who are aware of the early warning symptoms and signs of a Yes	895
Heart attack awareness: symptoms and importance of calling 9-1-1 (percent)	Percent of adults aged 20 years and over who are aware of the early warning symptoms and signs of a Yes	894
Heart attack Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have had a heart attack Yes Yes Yes Yes	283
Heart attack Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have had a heart attack Yes Yes Yes Yes	284
Heart attack patients given aspirin at hospital arrival (percent)	Hospital Compare: Heart attack patients given aspirin at hospital arrival Yes Yes Yes Yes	319
Heart attack patients prescribed ACE inhibitor or angiotensin receptor blocker at discharge (percent)	Hospital Compare: Heart attack patients prescribed angiotensin converting enzyme inhibitor or angiot Yes Yes Yes Yes	321
Heart attack patients with aspirin prescribed at hospital discharge (percent)	Hospital Compare: Heart attack patients with aspirin prescribed at hospital discharge Yes Yes Yes Yes	320
Heart attack patients with beta blocker prescribed at hospital discharge (percent)	Hospital Compare: Heart attack patients with beta blocker prescribed at hospital discharge Yes Yes Yes Yes	323
Heart attack patients with fibrinolytic received within 30 minutes of hospital arrival (percent)	Hospital Compare: Heart attack patients with fibrinolytic received within 30 minutes of hospital arrival Yes Yes Yes Yes	324
Heart attack patients with percutaneous coronary intervention within 90 minutes of hospital arrival (p	Hospital Compare: Heart attack patients with percutaneous coronary intervention within 90 minutes c Yes Yes Yes Yes	325
Heart attack patients with smoking cessation counseling during hospital stay (percent)	Hospital Compare: Heart attack patients with smoking cessation counseling during hospital stay Yes Yes Yes Yes	322
Heart disease deaths (per 100,000)	Heart disease death rate Yes Yes Yes	83
Heart disease prevalence: adults (percent)	Respondent-reported prevalence of heart disease among adults aged 18 and over Yes	50044
Heart failure hospitalizations: Medicare beneficiaries 65-74 years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 65 to 7 Yes	891
Heart failure hospitalizations: Medicare beneficiaries 75-84 years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 75 to 8 Yes	892
Heart failure hospitalizations: Medicare beneficiaries 85+ years (per 1,000)	Rate of hospitalizations of older adults with heart failure as the principal diagnosis-Adults aged 85 year Yes	893
Heart failure Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have heart failure Yes Yes Yes Yes	295
Heart failure Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have heart failure Yes Yes Yes Yes	296
Heart failure patients prescribed ACE inhibitor or angiotensin receptor blocker at discharge (percent)	Hospital Compare: Heart failure patients prescribed angiotensin converting enzyme inhibitor or angiot Yes Yes Yes Yes	328
Heart failure patients with discharge instructions (percent)	Hospital Compare: Heart failure patients with discharge instructions Yes Yes Yes Yes	326
Heart failure patients with evaluation of left ventricular systolic function (percent)	Hospital Compare: Heart failure patients with evaluation of left ventricular systolic function Yes Yes Yes Yes	327
Heart failure patients with smoking cessation counseling (percent)	Hospital Compare: Heart failure patients with smoking cessation counseling Yes Yes Yes Yes	329
Hepatitis A (per 100,000)	Hepatitis A rate Yes	967
Hepatitis A vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-2 doses hepatitis A v: Yes Yes	1000
Hepatitis A, acute (number)	Number of reported cases of acute Hepatitis A Yes Yes Yes	10022
Hepatitis B vaccination: health care workers (percent)	Percent of Hepatitis B vaccine coverage among high-risk populations-Health care personnel Yes	1016
Hepatitis B vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Hepatitis B vaccine (3 or more doses) Yes	1006
Hepatitis B, acute (number)	Number of reported cases of acute Hepatitis B Yes Yes Yes	10023
Hepatitis B: adults (per 100,000)	Rate of new hepatitis B infections in adults aged 19 and older Yes	988
Hepatitis B: children (per 100,000)	Rate of new hepatitis B cases among persons aged 2 to 18 years Yes	980
Hepatitis B: injection drug users (number)	Number of new hepatitis B infections among high-risk populations: Injection drug users Yes	989
Hepatitis B: men who have sex with men (number)	Number of new hepatitis B infections among high-risk populations: Men who have sex with men Yes	991
Hepatitis C (per 100,000)	Rate of new hepatitis C infections Yes	992
Hepatitis C infection status awareness (percent)	Percent of persons aware they have a hepatitis C infection Yes	1022
HepB vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-3 doses hepatitis B (t Yes Yes	995
Hib invasive disease: children <5 years (per 100,000)	Rate of serotype b cases of Haemophilus influenzae (Hib) invasive disease (children aged 5 years and u Yes	979
Hib vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-3 doses or more Haei Yes Yes	994
High blood cholesterol: adults with chronic kidney disease (percent)	Percent of adults with chronic kidney disease who have elevated lipid levels Yes	481
High blood pressure: adults 18+ (percent) (Source: BRFSS)	Percent of adults that report having been diagnosed with high blood pressure Yes Yes	123
High cholesterol Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have high cholesterol Yes Yes Yes Yes	50018
High cholesterol Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have high cholesterol Yes Yes Yes Yes	50023
High cholesterol: adults (percent)	Percent of adults with high total blood cholesterol levels Yes	885
High school completion: adults 18-24 years (percent)	Percent of persons aged 18-24 that completes high school education Yes	565
High school diploma receipt by students with disabilities (percent)	Percent of students who are served under the Individuals with Disabilities Education Act who graduate Yes	395
High school graduates, on-time with regular diploma (percent)	Percent of public school students who graduate with a regular diploma 4 years after starting 9th grade Yes	394

Hip fractures: female older adults (per 100,000)	Hip fracture hospitalizaion rate among females aged 65 years and over	Yes				429
Hip fractures: male older adults (per 100,000)	Hip fracture hospitalizaion rate among males aged 65 years and over	Yes				430
Hispanic Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who are Hispanic	Yes	Yes	Yes	Yes	313
HIV deaths (per 100,000)	HIV infection mortality rate	Yes	Yes	Yes		935
HIV infection status awareness (percent)	Percent of HIV-infected persons who are aware of their infection	Yes				942
HIV prevalence (per 100,000)	Rate of adolescents and adults aged 13 years and over living with HIV		Yes	Yes		10018
HIV testing: persons 15-24 years (percent)	Percent of adolescents and young adults who have been tested for HIV in the past 12 months	Yes				947
HIV testing: persons 15-44 years (percent)	Percent of adolescents and adults who have been tested for HIV in the past 12 months	Yes				944
HIV testing: pregnant women (percent)	Percent of pregnant women who have been tested for HIV during their pregnancy	Yes				946
HIV testing: TB patients (percent)	Percent of adults with tuberculosis (TB) who have been tested for HIV	Yes				934
HIV transmission rate: persons living with HIV (per 100)	Rate of HIV transmission per 100 persons with HIV among adolescents and adults	Yes				959
HIV/AIDS education: substance abuse treatment programs (percent)	Percent of substance abuse treatment facilities that offer HIV/AIDS education, counseling, and support	Yes				933
HIV/AIDS prevention discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about HIV/AIDS prevention before they we	Yes				814
HIV/AIDS prevention discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about HIV/AIDS prevention before th	Yes				815
HIV/AIDS prevention instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on HIV/AIDS prevention before they w	Yes				806
HIV/AIDS prevention instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on HIV/AIDS prevention before they we	Yes				807
Home Health Agencies (number)	Number of Home Health Agencies participating in the Medicare program	Yes	Yes		Yes	10041
Home health Medicare episodes (per 1,000 beneficiaries)	Average number of home health episodes among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2018
Home health Medicare users (count)	Medicare fee-for-service beneficiaries who used home health services	Yes	Yes	Yes	Yes	221
Home health Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used home health services	Yes	Yes	Yes	Yes	222
Home health Medicare visits (per 1,000 beneficiaries)	Average number of home health visits among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2017
Homes with dust-lead hazards (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US hom	Yes				723
Homes with lead paint hazards (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US hom	Yes				722
Homes with lead-based paint (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US hom	Yes				721
Homes with soil-lead hazards (number)	Number of U.S. homes that are found to have lead-based paint or related hazards: Number of US hom	Yes				724
Homicide deaths (per 100,000)	Homicide mortality rate	Yes	Yes	Yes		1041
Homicides, work-related (number)	Number of deaths due to work-related homicides	Yes				1299
Hospice Facilities (number)	Number of Hospice Facilities participating in the Medicare program	Yes	Yes		Yes	10042
Hospice Medicare admissions (per 1,000 beneficiaries)	Rate of hospice admissions among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2007
Hospice Medicare days (per 1,000 beneficiaries)	Average number of hospice days among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2015
Hospice Medicare users (count)	Medicare fee-for-service beneficiaries who had covered hospice services	Yes	Yes	Yes	Yes	227
Hospice Medicare utilization (percent)	Medicare fee-for-service beneficiaries who had covered hospice services	Yes	Yes	Yes	Yes	228
Hospital 30-day death (mortality) for heart attack patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for heart attack patients	Yes	Yes	Yes	Yes	343
Hospital 30-day death (mortality) for heart failure patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for heart failure patients	Yes	Yes	Yes	Yes	344
Hospital 30-day death (mortality) for pneumonia patients (percent)	Hospital Compare: Hospital 30-day death (mortality) rates for pneumonia patients	Yes	Yes	Yes	Yes	345
Hospital 30-day readmission for heart attack patients (percent)	Hospital Compare: Hospital 30-day readmission rates for heart attack patients	Yes	Yes	Yes	Yes	346
Hospital 30-day readmission for heart failure patients (percent)	Hospital Compare: Hospital 30-day readmission rates for heart failure patients	Yes	Yes	Yes	Yes	347
Hospital 30-day readmission for pneumonia patients (percent)	Hospital Compare: Hospital 30-day readmission rates for pneumonia patients	Yes	Yes	Yes	Yes	348
Hospital inpatient Medicare admissions (per 1,000 beneficiaries)	Rate of hospital inpatient admissions among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2001
Hospital inpatient Medicare days (per 1,000 beneficiaries)	Average number of hospital inpatient (IP) days among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2009
Hospital inpatient Medicare users (count)	Medicare fee-for-service beneficiaries who had covered hospital inpatient services	Yes	Yes	Yes	Yes	160
Hospital inpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries who had covered hospital inpatient services	Yes	Yes	Yes	Yes	161
Hospital outpatient Medicare users (count)	Medicare fee-for-service beneficiaries who used hospital outpatient services	Yes	Yes	Yes	Yes	187
Hospital outpatient Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used hospital outpatient services	Yes	Yes	Yes	Yes	188
Hospital outpatient Medicare visits (per 1,000 beneficiaries)	Rate of hospital outpatient visits among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2019
Housing units with physical problems (percent)	Percent of occupied housing units that have moderate or severe physical problems	Yes				663
How or why event occurred explained: news stories on foodborne disease outbreaks or national disast	Percent of crisis and emergency risk messages embedded in print and broadcast news stories that expl	Yes				10096
HPV vaccination: girls 13-15 years (percent)	Percent of female adolescents who have 3 doses Human papillomavirus vaccine (HPV) by age 13 to 15	Yes	Yes			1013

Hypertension admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Hypertension admission rate among Medicare fee-for-service beneficia	Yes	Yes	Yes	Yes	2040
Hypertension Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have hypertension	Yes	Yes	Yes	Yes	50009
Hypertension Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have hypertension	Yes	Yes	Yes	Yes	50010
Hypertension medication compliance (percent)	Percent of adults with hypertension who are taking the prescribed medications to lower their blood pr	Yes				912
Hypertension: adults (percent)	Percent of adults with hypertension	Yes				897
Hypertension: adults with chronic kidney disease (percent)	Percent of adults with chronic kidney disease who have elevated blood pressure	Yes				480
Hypertension: children 8-17 years (percent)	Percent of children and adolescents with hypertension	Yes				898
Identification of "people with disabilities" in data systems (number)	Number of major applicable Healthy People 2020 data systems with a standardized set of questions th	Yes				513
Illegal drugs on school property (percent)	Percent of adolescents who have been offered, sold, or given an illegal drug on school property	Yes	Yes			411
Illicit drug abstinence, prenatal (percent)	Abstinence from alcohol, cigarettes, and illicit drugs among pregnant women-Illicit drugs	Yes				1145
Illicit drug treatment receipt (percent)	Percent of persons who need illicit drug treatment and received specialty treatment for abuse or depe	Yes				1454
Illicit drugs, never use: high school seniors (percent)	Percent of high school seniors never using substances-Illicit drugs	Yes				1431
Illicit drugs, recent use: adults (percent)	Percent of adults reporting use of any illicit drug during the past 30 days	Yes				1434
Illicit drugs, recent use: teens (percent)	Percent of adolescents reporting use of alcohol or any illicit drugs during the past 30 days	Yes				1432
Imaging Medicare service events (per 1,000 beneficiaries)	Rate of imaging service events among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2023
Imaging Medicare users (count)	Medicare fee-for-service beneficiaries who used imaging services	Yes	Yes	Yes	Yes	251
Imaging Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used imaging services	Yes	Yes	Yes	Yes	252
Immunization information system participation: States (number)	Number of States that have 80 percent of adolescents with two or more age-appropriate immunizatio	Yes				1025
Impaired fecundity: women 18-44 years (percent)	Percent of women aged 18 to 44 years who have impaired fecundity	Yes				1193
Improvement plan development: health emergencies/exercises (days)	Time reported for State public health agencies to establish improvement plans following responses to	Yes				1397
Inability to obtain or delay in obtaining dental care (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary dental care	Yes				381
Inability to obtain or delay in obtaining medical care (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary medical care	Yes				380
Inability to obtain or delay in obtaining prescription meds (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary prescription medicines	Yes				382
Inability to obtain or delayed access to medical/dental care or prescriptions (percent)	Percent of individuals who are unable to obtain or delay in obtaining necessary medical care, dental ca	Yes				379
Incentive policies for food retail: states (number)	Number of states and the District of Columbia that have state-level policies that incentivize food retail	Yes				1239
Inclusion in regular education: students with disabilities (percent)	Percent of children and youth with disabilities in classroom settings who spend at least 80 percent of t	Yes				517
Increasing physical activity: adults with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report increasing their levels of ph	Yes				562
Indoor allergen levels, cockroach (units of allergen/gram of settled dust)	Indoor allergen levels: cockroach	Yes				679
Indoor allergen levels, mouse (micrograms of allergen/gram of settled dust)	Indoor allergen levels: mouse	Yes				680
Infant deaths between 28 days-1 year (per 1,000 live births)	Rate of postneonatal deaths (between 28 days and 1 year)	Yes	Yes	Yes		1160
Infant deaths due to birth defects (per 1,000 live births)	Rate of infant deaths related to birth defects (all birth defects)	Yes				1161
Infant deaths due to congenital heart defects (per 1,000 live births)	Rate of infant deaths related to birth defects (congenital heart defects)	Yes				1162
Infant deaths from sudden infant death syndrome (per 1,000 live births)	Rate of infant deaths from sudden infant death syndrome (SIDS)	Yes				1163
Infant deaths from sudden unexpected/unexplained causes (per 1,000 live births)	Rate of infant deaths from sudden unexpected/unexplained causes (includes SIDS, Unknown Cause, Ac	Yes				1164
Infant deaths within first 28 days of life (per 1,000 live births)	Rate of neonatal deaths (within the first 28 days of life)	Yes	Yes	Yes		1159
Infant deaths, all (per 1,000 live births)	Rate of all infant deaths (within 1 year)	Yes	Yes	Yes		1158
Infants put to sleep on their backs (percent)	Percent of infants who are put down to sleep on their backs	Yes				1141
Influenza and pneumonia deaths (per 100,000)	Influenza and pneumonia death rate per 100,000	Yes	Yes	Yes		50013
Influenza-virus resistance to antivirals monitoring: states (number)	Number of public health laboratories monitoring influenza-virus resistance to antiviral agents	Yes				1028
Information systems used for environmental health: states (number)	Quality, utility, awareness, and use of existing information systems for environmental health among st	Yes				664
Inhalant use: teens (percent)	Percent of adolescents who use inhalants	Yes				1443
Injection drug use admissions (number)	Number of admissions to substance abuse treatment for injection drug use	Yes				1453
Injuries due to falls: adults with balance/dizziness problems (percent)	Percent of adults with balance and dizziness problems who have been injured as a result of a fall for ar	Yes				770
Injuries, nonfatal unintentional (per 100,000)	Nonfatal unintentional injury rate	Yes				1075
Injuries, non-fatal, work-related (per 100)	Rate of injuries in private sector industries resulting in medical treatment, lost time from work, or rest	Yes				1308
Injuries, sports and recreation (per 1,000)	Sports and recreation injury rate	Yes				1098
Injuries, treated in ED, non-fatal, work-related (per 100)	Rate of injuries treated in emergency departments from work, or restricted work activity, as reported	Yes				1309
Injuries, treated in ED, non-fatal, work-related: teens (per 100)	Rate of nonfatal work-related injuries among dolescent workers aged 15 to 19 years	Yes				1310

Injury deaths (per 100,000)	Injury death rate	Yes	Yes	Yes	1071
Injury deaths, unintentional (per 100,000)	Unintentional injury death rate	Yes	Yes	Yes	1074
Injury deaths, work-related, all industries (per 100,000)	Rate of work-related injury deaths - All industry	Yes			1303
Injury deaths, work-related, construction (per 100,000)	Rate of work-related injury deaths - Construction	Yes			1305
Injury deaths, work-related, farming/forestry/fishing/hunting (per 100,000)	Rate of work-related injury deaths - Agriculture, forestry, fishing, and hunting	Yes			1307
Injury deaths, work-related, mining (per 100,000)	Rate of work-related injury deaths - Mining	Yes			1304
Injury deaths, work-related, transportation and warehousing: persons 16+ years (per 100,000)	Rate of work-related injury deaths - Transportation and warehousing: persons 16+ years	Yes			1306
Injury ED visits (per 100,000)	Rate of emergency department visits for nonfatal injuries	Yes			1073
Injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal injuries	Yes			1072
Injury prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, Yes	Yes			596
Inpatient rehab Medicare users (count)	Medicare fee-for-service beneficiaries who had covered inpatient rehabilitation facility services	Yes	Yes	Yes	207
Inpatient rehab Medicare utilization (percent)	Medicare fee-for-service beneficiaries who had covered inpatient rehabilitation facility services	Yes	Yes	Yes	208
Inpatient rehabilitation facility Medicare admissions (per 1,000 beneficiaries)	Rate of inpatient rehabilitation facility (IRF) admissions among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	2005
Inpatient rehabilitation facility Medicare days (per 1,000 beneficiaries)	Average number of inpatient rehabilitation facility (IRF) days among Medicare fee-for-service benefici	Yes	Yes	Yes	2013
Inspection of drinking water for bacteria: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes			690
Inspection of drinking water for coliforms: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes			691
Inspection of drinking water for lead: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes			689
Intended pregnancies (percent)	Percent of pregnancies that are intended	Yes			779
Internet access (percent)	Percent of persons with access to the Internet	Yes			857
Internet use to communicate with providers (percent)	Percent of persons who use the Internet to communicate with their provider	Yes			868
Internet use to track personal health information (percent)	Percent of persons who use the Internet to keep track of personal health information, such as care rec	Yes			867
Interprofessional educational experiences: schools with health professions education programs (percent)	Percent of academic institutions with health professions education programs whose prevention curricula	Yes			10087
Invasive colorectal cancer (per 100,000)	Invasive colorectal cancer incidence	Yes			509
Invasive pneumococcal infections: children <5 years (per 100,000)	Rate of new invasive pneumococcal infections among children under age 5 years	Yes			963
Invasive pneumococcal infections: older adults (per 100,000)	Rate of new invasive pneumococcal infections among adults aged 65 years and older	Yes			964
Invasive uterine cervical cancer: females (per 100,000)	Invasive uterine cervical cancer incidence	Yes			510
Involvement in health care decision making (percent)	Percent of persons who report that their health care providers always involved them in decisions about	Yes			866
Iron deficiency: children 1-2 years (percent)	Percent of young children aged 1 to 2 years who are iron deficient	Yes			1209
Iron deficiency: children 3-4 years (percent)	Percent of young children aged 3 to 4 years who are iron deficient	Yes			1210
Iron deficiency: females of childbearing age (percent)	Percent of females of childbearing age, aged 12 to 49 years, who are iron deficient	Yes			1211
Iron deficiency: pregnant females (percent)	Percent of pregnant females who are iron deficient	Yes			1212
Ischemic heart disease Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have ischemic heart disease	Yes	Yes	Yes	297
Ischemic heart disease Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have ischemic heart disease	Yes	Yes	Yes	298
Joint difficulty grasping: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-re	Yes			434
Joint difficulty stooping/bending/ kneeling: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-re	Yes			433
Joint difficulty walking 3 blocks: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-re	Yes			431
Joint difficulty walking up 10 steps: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who find it "very difficult" to perform specific joint-re	Yes			432
Joint pain: adults with arthritis (mean)	Mean level of joint pain among adults with doctor-diagnosed arthritis	Yes			418
Kidney diseases deaths (per 100,000)	Nephritis, nephrotic syndrome, and nephrosis (kidney diseases) death rate per 100,000	Yes	Yes	Yes	50014
Kidney transplantation within 3 years: hemodialysis patients (percent)	Percent of patients receiving a kidney transplant within 3 years of end-stage renal disease (ESRD)	Yes			473
Kidney waitlisting/transplantation: hemodialysis patients < 70 years (percent)	Percent of hemodialysis patients wait-listed and/or receiving a deceased donor kidney transplant with	Yes			472
Kindergarten vaccination record keeping: states (number)	Number of States collecting kindergarten vaccination coverage data according to CDC minimum stand	Yes			1026
Known health risks explained: news stories on foodborne disease outbreaks or national disasters (percent)	Percent of crisis and emergency risk messages embedded in print and broadcast news stories that expl	Yes			10094
Laboratories support data management: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory ser	Yes			1350
Laboratories support disease prevention/control: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory ser	Yes			1349
Laboratories support emergency response: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory ser	Yes			1356
Laboratories support environmental health: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory ser	Yes			1352
Laboratories support food safety: States (percent)	Percent of Tribal and State public health agencies that provide or assure comprehensive laboratory ser	Yes			1353

Mandatory ignition interlock laws: States (number)	Number of States with mandatory ignition interlock laws for first and repeat impaired driving offender	Yes				1468
Marijuana, never use: at risk teens (percent)	Percent of at risk adolescents aged 12 to 17 years who, in the past year, refrained from using marijuana	Yes				1429
Marijuana, recent use: teens (percent)	Percent of adolescents reporting use of marijuana during the past 30 days	Yes				1433
Mass transit for transportation (percent)	Percent of trips to work using of alternative modes of transportation - mass transit	Yes				668
Maternal deaths (per 100,000 live births)	Rate of maternal death	Yes				1129
Maternal illness and complications due to pregnancy (percent)	Maternal illness and complications due to pregnancy (complications during hospitalized labor and deli	Yes				1130
Measles, total (number)	Number of reported cases of measles	Yes	Yes	Yes		10032
Measles, U.S.-acquired (number)	Number of measles cases (U.S.-acquired cases)	Yes				981
Median Household Income (dollars)	Small area estimates of median household income	Yes	Yes	Yes		10003
Medicaid beneficiaries (number in thousands)	Number of Medicaid beneficiaries	Yes	Yes			10014
Medicaid coverage for nicotine treatment: States (number)	Comprehensive Medicaid insurance coverage of evidence-based treatment for nicotine dependency in	Yes				1516
Medicaid coverage: persons <65 (percent)	Percent of persons under age 65 who have Medicaid coverage	Yes				50057
Medicaid managed care beneficiaries (percent)	Percent of Medicaid beneficiaries in managed care plans	Yes	Yes			10015
Medicaid payments per beneficiary (dollars)	Average payment per Medicaid beneficiary	Yes	Yes			10017
Medical care delayed or not received due to cost (percent) (Source: NHIS)	Percent of persons who did not get or delayed medical care due to cost in the previous 12 months	Yes	Yes			50065
Medical evaluation: older adults with chronic kidney disease & diabetes (percent)	Percent of persons 65 years of age and over with type 1 or type 2 diabetes and chronic kidney disease	Yes				478
Medical evaluation: older adults with chronic kidney disease (percent)	Percent of persons 65 years of age and over with chronic kidney disease who receive medical evaluati	Yes				477
Medical home: children (percent)	Percent of children who have access to a medical home	Yes				1152
Medical home: children with special needs (percent)	Percent of children with special health care needs who have access to a medical home	Yes				1153
Medical treatment: older adults with chronic kidney disease & diabetes (percent)	Percent of persons 65 years of age and over with Part D insurance with diabetes and chronic kidney di	Yes				479
Medicare beneficiaries who are also eligible for Medicaid (percent)	Medicare fee-for-service beneficiaries who are also eligible for Medicaid	Yes	Yes	Yes	Yes	317
Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	307
Medicare managed care beneficiaries (percent)	Percent of Medicare beneficiaries enrolled in Medicare Advantage managed care plans	Yes	Yes			10020
Melanoma cancer deaths (per 100,000)	Melanoma cancer death rate	Yes				493
Meningococcal disease (number)	Meningococcal disease rate	Yes				968
Meningococcal vaccination: teens 13-15 years (percent)	Percent of adolescents who have 1 dose Meningococcal vaccine (MCV) by age 13 to 15 years	Yes	Yes			1012
Mental health screening: juvenile justice facilities (percent)	Percent of juvenile residential facilities that screen admissions for mental health problems	Yes				1111
Mental health services receipt: homeless adults (percent)	Percent of homeless adults with mental health problems who receive mental health services	Yes				1107
Mental health treatment provision: primary care facilities (percent)	Percent of primary care facilities that provide mental health treatment onsite or by paid referral	Yes				1109
Mental illness prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services	Yes				598
Mentally unhealthy days: adults (per person)	Average number of reported mentally unhealthy days per month among adults 18 years and over		Yes	Yes		9
Mercury level: children 1-5 years (µg/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				698
Mercury level: females 16-49 years (µg/L)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				699
Mercury poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that ca	Yes				715
Methicillin-resistant Staphylococcus aureus infections (per 100,000)	Invasive healthcare-associated methicillin-resistant Staphylococcus aureus (MRSA) infections	Yes				852
Missed school days: children 5-17 with asthma (percent)	Percent of children aged 5 to 17 years with asthma who miss school days	Yes				1410
Missed work days: adults 18-64 with asthma (percent)	Percent of adults aged 18 to 64 years with asthma who miss work days	Yes				1411
MMR vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-1 dose measles-mum	Yes	Yes			996
MMR vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-MMR vaccines (2 or more doses)	Yes				1004
Mold problem policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes				684
Motor vehicle crash deaths (per 100,000,000 vehicle miles traveled)	Motor vehicle crash-related deaths per 100 million vehicle miles traveled	Yes				1077
Motor vehicle crashes involving drowsy driving (per 100,000,000 vehicle miles traveled)	Motor vehicle crashes per 100 million miles traveled involving drowsy driving	Yes				1470
Motor vehicle injuries, nonfatal (per 100,000)	Nonfatal motor vehicle crash-related injuries	Yes				1078
Motor vehicle traffic deaths (per 100,000)	Motor vehicle traffic-related death rate per 100,000	Yes	Yes	Yes		1076
Motorcycle helmet use (percent)	Percent of motorcycle operators and passengers using helmets	Yes				1048
Multivitamins/folic acid use, preconception (percent)	Percent of women delivering a live birth who took multivitamins/folic acid prior to pregnancy	Yes				1179
Mumps (number)	Number of reported cases of mumps	Yes	Yes	Yes		10031
Mumps, U.S.-acquired (number)	Number of mumps cases (U.S.-acquired cases)	Yes				982

Muscle-strengthening activity meeting guidelines: adults (percent)	Percent of adults who perform muscle-strengthening activities on 2 or more days of the week	Yes				1321
Neck pain: adults (percent)	Percent of adults aged 18 and over with neck pain during the past three months	Yes				50041
Neonatal hearing screening (percent)	Percent of infants that receive screening for hearing loss no later than age 1 month	Yes				756
Nephrologist seen at least 12 months before hemodialysis initiation (percent)	Percent of chronic kidney disease patients receiving care from a nephrologist at least 12 months before	Yes				468
Never had sexual intercourse: females <16 years (percent)	Percent of female adolescents aged 15 years and under who have never had sexual intercourse	Yes				792
Never had sexual intercourse: females 15-17 years (percent)	Percent of female adolescents aged 15 to 17 years who have never had sexual intercourse	Yes				790
Never had sexual intercourse: males <16 years (percent)	Percent of male adolescents aged 15 years and under who have never had sexual intercourse	Yes				793
Never had sexual intercourse: males 15-17 years (percent)	Percent of male adolescents aged 15 to 17 years who have never had sexual intercourse	Yes				791
Newborn screening verification: states (number)	Number of States and the District of Columbia that verify through linkage with vital records that all ne	Yes				1173
No exercise: adults (percent) (Source: BRFSS)	Adults that report no leisure-time exercise in past month		Yes	Yes		120
No health care visits: children (percent)	Percent of children under age 18 years who had no health care visits to an office or clinic within the pa	Yes				50068
Nonfatal firearm-related injuries (per 100,000)	Nonfatal firearm-related injury rate	Yes				1043
Non-FDA-approved pain medications (number)	Number of non-FDA-approved pain medications	Yes				1198
Non-Hispanic White Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who are Non-Hispanic white	Yes	Yes	Yes	Yes	311
Number of days the Air Quality Index (AQI) exceeds 100 (days)	Reduce the number of days the Air Quality Index (AQI) exceeds 100, weighted by population and AQI v	Yes				665
Nurse practitioner training, cultural diversity (percent)	Percent of nurse practitioner programs that include training in cultural diversity	Yes				625
Nurse practitioner training, environmental health (percent)	Percent of nurse practitioner programs that include training in environmental health	Yes				627
Nurse practitioner training, evaluation of health literature (percent)	Percent of nurse practitioner programs that include training in evaluation of health sciences literature	Yes				626
Nurse practitioner training, global health (percent)	Percent of nurse practitioner programs that include training in global health	Yes				629
Nurse practitioner training, health promotion counseling (percent)	Percent of nurse practitioner programs that include training in counseling for health promotion and di	Yes				624
Nurse practitioner training, public health systems (percent)	Percent of nurse practitioner programs that include training in public health systems	Yes				628
Nursing home (Title 18/19) facilities (number)	Number of Medicare or Medicaid-certified Title 18/19 (nursing home) facilities	Yes	Yes		Yes	10053
Nursing training, counseling for health promotion (percent)	Percent of undergraduate nursing programs that include training in counseling for health promotion a	Yes				618
Nursing training, cultural diversity (percent)	Percent of undergraduate nursing programs that include training in cultural diversity	Yes				619
Nursing training, environmental health (percent)	Percent of undergraduate nursing programs that include training in environmental health	Yes				621
Nursing training, evaluation of health sciences literature (percent)	Percent of undergraduate nursing programs that include training in evaluation of health sciences litera	Yes				620
Nursing training, global health (percent)	Percent of undergraduate nursing programs that include training in global health	Yes				623
Nursing training, public health systems (percent)	Percent of undergraduate nursing programs that include training in public health systems	Yes				622
Nutrition counseling at physician visits (percent)	Percent of physician visits made by all child or adult patients that include counseling about nutrition o	Yes				1227
Nutrition counseling: patients with CVD, diabetes, or hyperlipidemia (percent)	Percent of physician office visits made by patients with a diagnosis of cardiovascular disease, diabetes, Yes	Yes				1225
Nutrition services: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services, Yes	Yes				603
Nutrition standards for child care: states (number)	Number of states and the District of Columbia with nutrition standards for foods and beverages provid	Yes				1240
Obesity: adults (percent)	Percent of adults who are obese	Yes				1208
Obesity: adults (percent) (Source: BRFSS)	Percent of adults 18 years and over that report BMI >= 30		Yes	Yes		15
Obesity: children 2-19 years (percent)	Percent of children and adolescents aged 2 to 19 years who are considered obese	Yes				1216
Obesity: children 2-5 years (percent)	Percent of children and adolescents aged 2 to 5 years who are considered obese	Yes				1213
Obesity: children 6-11 years (percent)	Percent of children and adolescents aged 6 to 11 years who are considered obese	Yes				1214
Obesity: teens 12-19 years (percent)	Percent of children and adolescents aged 12 to 19 years who are considered obese	Yes				1215
Obstructive sleep apnea evaluation: adults (percent)	Percent of adults with symptoms of obstructive sleep apnea who seek medical evaluation	Yes				1469
Occupational eye injuries resulting in lost work days (per 10,000)	Rate of occupational eye injuries resulting in lost work days	Yes				1564
Occupational eye injuries treated in emergency departments (per 10,000)	Rate of occupational eye injuries treated in emergency departments	Yes				1565
Online health information access (percent)	Percent of online health information seekers who report easily accessing health information	Yes				865
Oral and craniofacial health surveillance: states (number)	Number of States and the District of Columbia that have an oral and craniofacial health surveillance sy	Yes				1268
Oral and pharyngeal cancer early detection (percent)	Percent of oral and pharyngeal cancers detected at the earliest stage	Yes				1264
Oral health care system use (percent)	Percent of children, adolescents, and adults who used the oral health care system in the past year	Yes				1266
Oral health program: Federally qualified health centers (percent)	Percent of Federally Qualified Health Centers that have an oral health care program	Yes				1287
Oral health program: Local health departments (percent)	Percent of local health departments that have oral health prevention or care programs	Yes				1288
Oral health services receipt at health centers (percent)	Percent of patients who receive oral health services at Federally Qualified Health Centers each year	Yes				1296

Organ Procurement Organizations (number)	Number of Organ Procurement Organizations participating in the Medicare program	Yes	Yes	Yes	10044
Oropharyngeal cancer deaths (per 100,000)	Oropharyngeal cancer death rate	Yes			491
Osteoporosis Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have osteoporosis	Yes	Yes	Yes	50019
Osteoporosis Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have osteoporosis	Yes	Yes	Yes	50024
Osteoporosis: adults 50+ years (percent)	Percent of adults aged 50 years and over with osteoporosis	Yes			427
Other or Unknown race Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who are Other or Unknown race	Yes	Yes	Yes	316
Otitis media visits in ambulatory care settings: children (per 1,000)	Rate of otitis media in children and adolescents	Yes			743
Outbreak-associated infections associated with beef (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacte	Yes			839
Outbreak-associated infections associated with chicken (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacte	Yes			843
Outbreak-associated infections associated with dairy (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacte	Yes			840
Outbreak-associated infections associated with fruits/nuts (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacte	Yes			841
Outbreak-associated infections associated with vegetables (number)	Number of outbreak-associated infections due to Shiga toxin-producing E. coli O157, or Campylobacte	Yes			842
Outpatient physical therapy/speech pathology facilities (number)	Number of Outpatient physical therapy/speech pathology facilities participating in the Medicare progr	Yes	Yes	Yes	10045
Overdose ED visits for injectable antidiabetic agents (per 10,000 outpatient prescription visits)	Rate of emergency department (ED) visits for overdoses from injectable antidiabetic agents	Yes			1204
Overdose ED visits for narrow-therapeutic-index medications, (per 10,000 outpatient prescription visit	Rate of emergency department (ED) visits for overdoses of narrow-therapeutic-index medications	Yes			1205
Overdose ED visits for oral anticoagulants (per 10,000 outpatient prescription visits)	Rate of emergency department (ED) visits for overdoses from oral anticoagulants	Yes			1203
Overdose ED visits: children <5 years (per 10,000)	Rate of emergency department (ED) visits for medication overdoses among children less than 5 years c	Yes			1206
Overexertion or repetitive motion injuries/illness from work (per 10,000)	Rate of injury and illness cases involving days away from work due to overexertion or repetitive motio	Yes			1297
Ozone days (number)	Annual number of days with daily 8-hour maximum ozone concentration over the National Ambient Ai			Yes	45
Pap smears: women 18+ (percent) (Source: NHIS)	Percent of women aged 18 and over who have had a Pap smear within the past 3 years	Yes			50052
Pap smears: women 18+ without hysterectomy (percent)	Percent of women aged 18 and over who have not had a hysterectomy who have had a Pap smear wit	Yes			50059
Pap test counseling: women 21-65 years (percent)	Percent of women who were counseled by their providers about Pap tests	Yes			495
Pap test: women 18+ (percent)	Percent of women age 18+ that report Pap test in the last 3 years		Yes	Yes	115
Para-nitrophenol level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes			703
Parent employed year round and full time: children (percent)	Percent of children aged 0-17 years living with at least one parent employed year round, full time	Yes			10070
Part B drug Medicare users (count)	Medicare fee-for-service beneficiaries who used Part B drugs	Yes	Yes	Yes	269
Part B drugs Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used Part B drugs	Yes	Yes	Yes	270
Particulate matter days (percent)	Percent of days with maximum 24-hour average PM 2.5 concentration over the National Ambient Air C			Yes	44
PCB level, dioxin-like: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes			707
PCB level, nondioxin-like: persons 12+ years (ng/g of lipid)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes			706
PCV vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-4 doses pneumococc	Yes	Yes		999
Pedal-cyclist deaths (per 100,000)	Death rate of pedal-cyclists on public roads	Yes			1103
Pedestrian deaths (per 100,000)	Rate of pedestrian deaths on public roads	Yes			1045
Pedestrian injuries, nonfatal (per 100,000)	Rate of nonfatal pedestrian injuries on public roads	Yes			1046
Pelvic inflammatory disease (percent)	Percent of females aged 15 to 44 years who have ever required treatment for pelvic inflammatory dise	Yes			1473
Perchlorate level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes			710
Percutaneous intervention timeliness: heart attack patients (percent)	Percent of eligible patients with heart attacks who receive Percutaneous intervention (PCI) within 90 n	Yes			879
Perinatal deaths (per 1,000 live births plus fetal deaths)	Rate of fetal and infant deaths during perinatal period (28 weeks of gestation to 7 days after birth)	Yes			1157
Periodontal disease: adults 45-74 years (percent)	Percent of adults aged 45 to 74 years with periodontitis	Yes			1280
Personal care limitations: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who have difficulty in performing two or more perso	Yes			420
Pertussis (number)	Number of reported cases of pertussis	Yes	Yes	Yes	10029
Pertussis: infants (number)	Number of pertussis cases among children under age 1 year	Yes			983
Pertussis: teens (number)	Number of pertussis cases among adolescents aged 11 to 18 years	Yes			984
Pesticide area marking policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes			687
Pesticide exposures causing health care facility visits (number)	Number of pesticide exposures that result in visits to a health care facility	Yes			661
Pesticide information policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes			688
Pesticide limited use policy: schools (percent)	Percent of the Nation's elementary, middle, and high schools that have official school policies and eng	Yes			686
Pesticide poisoning monitoring: states (number)	Number of States, Territories, and the District of Columbia that monitor diseases or conditions that ca	Yes			714

Pharmacist training, public health systems (percent)	Percent of Doctor of Pharmacy (PharmD) granting colleges and schools of pharmacy that include traini	Yes				10079
Pharmacist training, counseling for health promotion (percent)	Percent of Doctor of Pharmacy (PharmD) granting colleges and schools of pharmacy that include traini	Yes				10075
Pharmacist training, cultural diversity (percent)	Percent of Doctor of Pharmacy (PharmD) granting colleges and schools of pharmacy that include traini	Yes				10076
Pharmacist training, environmental health (percent)	Percent of Doctor of Pharmacy (PharmD) granting colleges and schools of pharmacy that include traini	Yes				10078
Pharmacist training, evaluation of health sciences literature (percent)	Percent of Doctor of Pharmacy (PharmD) granting colleges and schools of pharmacy that include traini	Yes				10077
Pharmacist training, global health (percent)	Percent of Doctor of Pharmacy (PharmD) granting colleges and schools of pharmacy that include traini	Yes				10080
Phthalate level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				711
Physical activity counseling: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who receive health care provider counseling-For phy	Yes				422
Physical activity duration policies for child care settings: States (number)	Number of States with licensing regulations for physical activity provided in child care-Require number	Yes				1345
Physical activity intensity policies for child care settings: States (number)	Number of States with licensing regulations for physical activity provided in child care-Require childre	Yes				1344
Physical activity policies for child care settings: States (number)	Number of States with licensing regulations for physical activity provided in child care-Require activity	Yes				1343
Physical activity: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services,	Yes				604
Physical assault injuries, nonfatal (per 100,000)	Nonfatal physical assault injury rate	Yes				1099
Physical assaults: persons 12+ years (per 1,000)	Rate of physical assaults among persons 12 years of age and over	Yes				1051
Physical education required daily: elementary schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students	Yes				1314
Physical education required daily: high schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students	Yes				1316
Physical education required daily: middle/junior high schools (percent)	Percent of the Nation's public and private schools that require daily physical education for all students	Yes				1315
Physical education, daily at school: teens (percent)	Percent of adolescents who participate in daily school physical education	Yes	Yes			1317
Physical fighting: teens (percent)	Percent of physical fighting among adolescents	Yes	Yes			1052
Physically or mentally unhealthy days: adults (days)	Average number of reported physically or mentally unhealthy days per month among persons 18 year:	Yes	Yes			75
Physically unhealthy days: adults (per person)	Average number of reported physically unhealthy days per month among adults 18 years of age and o	Yes	Yes			7
Physician asst. training in cultural diversity (percent)	Percent of physician assistant programs that include training in cultural diversity	Yes				631
Physician asst. training, counseling for health promotion (percent)	Percent of physician assistant programs that include training in counseling for health promotion and d	Yes				630
Physician asst. training, environmental health (percent)	Percent of physician assistant programs that include training in environmental health	Yes				633
Physician asst. training, evaluation of health literature (percent)	Percent of physician assistant programs that include training in evaluation of health sciences literature	Yes				632
Physician asst. training, global health (percent)	Percent of physician assistant programs that include training in global health	Yes				635
Physician asst. training, public health systems (percent)	Percent of physician assistant programs that include training in public health systems	Yes				634
Physician evaluation and management Medicare service events (per 1,000 beneficiaries)	Rate of physician evaluation and management service events among Medicare fee-for-service benefi	Yes	Yes	Yes	Yes	2021
Physician evaluation and management Medicare users (count)	Medicare fee-for-service beneficiaries who used physician evaluation and management services	Yes	Yes	Yes	Yes	239
Physician evaluation and management Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used physician evaluation and management services	Yes	Yes	Yes	Yes	240
Physician physical activity counseling (percent)	Percent of physician visits made by all child and adult patients that include counseling about exercise	Yes				1342
Physician physical activity counseling: adults with chronic conditions (percent)	Percent of office visits made by patients with a diagnosis of cardiovascular disease, diabetes, or hyperl	Yes				1341
Physician procedures Medicare service events (per 1,000 beneficiaries)	Rate of physician procedure service events among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2022
Physician procedures Medicare users (count)	Medicare fee-for-service beneficiaries who used physician procedures services	Yes	Yes	Yes	Yes	245
Physician procedures Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used physician procedures services	Yes	Yes	Yes	Yes	246
Physician use delayed due to cost (percent) (Source: BRFS)	Percent of adults 18+ who needed to see a doctor but could not because of cost	Yes	Yes			10019
Pneumococcal vaccination: adults 65+ (percent) (Source: BRFS)	Percent of adults age 65+ that report ever having a pneumococcal vaccination	Yes	Yes			118
Pneumococcal vaccination: high-risk adults (percent)	Percent of adults who are vaccinated against pneumococcal disease-Noninstitutionalized high-risk adu	Yes				1019
Pneumococcal vaccination: nursing home residents (percent)	Percent of adults who are vaccinated against pneumococcal disease-Institutionalized adults (persons a	Yes				1020
Pneumococcal vaccination: older adults (percent)	Percent of adults who are vaccinated against pneumococcal disease-Noninstitutionalized adults aged 6	Yes				1018
Pneumoconiosis deaths (number)	Annual number of pneumoconiosis deaths	Yes				1298
Pneumonia patient with blood cultures in emergency department before antibiotic administered (perc	Hospital Compare: Pneumonia patient with blood cultures in emergency department before antibiotic	Yes	Yes	Yes	Yes	332
Pneumonia patients with appropriate initial antibiotic selection for community-acquired pneumonia (p	Hospital Compare: Pneumonia patients with appropriate initial antibiotic selection for community-acq	Yes	Yes	Yes	Yes	331
Pneumonia patients with influenza vaccination (percent)	Hospital Compare: Pneumonia patients with influenza vaccination	Yes	Yes	Yes	Yes	333
Pneumonia patients with initial antibiotic received within 6 hours of hospital arrival (percent)	Hospital Compare: Pneumonia patients with initial antibiotic received within 6 hours of hospital arrival	Yes	Yes	Yes	Yes	335
Pneumonia patients with pneumococcal vaccination (percent)	Hospital Compare: Pneumonia patients with pneumococcal vaccination	Yes	Yes	Yes	Yes	330
Pneumonia patients with smoking cessation counseling (percent)	Hospital Compare: Pneumonia patients with smoking cessation counseling	Yes	Yes	Yes	Yes	334
Poisoning deaths (per 100,000)	Poisoning death rate: All persons	Yes	Yes	Yes		1064

Poisoning deaths, unintentional/undetermined (per 100,000)	Poisoning death rate of unintentional or undetermined intent among all persons	Yes			1066
Poisoning deaths, unintentional/undetermined: adults 35-54 years (per 100,000)	Poisoning death rate of unintentional or undetermined intent among persons aged 35 to 54 years	Yes			1067
Poisoning deaths: adults 35-54 years (per 100,000)	Poisoning death rate: Persons aged 35 to 54 years	Yes			1065
Poisonings, nonfatal (per 100,000)	Rate of nonfatal poisonings	Yes			1040
Polio vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-3 doses polio vaccine	Yes	Yes		997
Polio vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Polio vaccine (3 or more doses)	Yes			1005
Polio, U.S.-acquired (number)	Number of acute paralytic poliomyelitis cases (U.S.-acquired cases)	Yes			985
Population density (per square mile)	Population density of United States geographic area per square mile	Yes	Yes	Yes	10001
Population distributions (percent)	Bridged-race population distributions according to 1977 OMB standard	Yes	Yes	Yes	10002
Population estimates (number)	Bridged-race population estimates according to 1977 OMB standard	Yes	Yes	Yes	10000
Population-based Healthy People 2020 objectives with race and ethnicity data (percent)	Percent of population-based Healthy People 2020 objectives for which national data are available by r	Yes			10091
Population-based Healthy People 2020 objectives with sex data (percent)	Percent of population-based Healthy People 2020 objectives for which national data are available by s	Yes			10092
Population-based Healthy People 2020 objectives with socioeconomic status data (percent)	Percent of population-based Healthy People 2020 objectives for which national data are available by s	Yes			10093
Portable X-Ray providers (number)	Number of portable X-Ray providers participating in the Medicare Program	Yes	Yes	Yes	10046
Positive communication with children: parents (percent)	Percent of parents who use positive communication with their child	Yes			739
Postdiarrheal hemolytic-uremic syndrome infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Postdiarrheal hemol	Yes			826
Poverty (percent)	Percent of persons of all ages living in poverty	Yes			10072
Poverty: children < 18 years (percent)	Percent of children aged 0-17 years living in poverty	Yes			10073
Poverty: children < 18 years (percent)	Percent of children under 18 years of age living in poverty	Yes	Yes	Yes	36
Preemptive smoke-free air laws: States (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption on sm	Yes			1545
Preemptive tobacco advertising laws: States (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption in adv	Yes			1546
Preemptive tobacco control laws, youth access: States (number)	Number of States with State laws that preempt stronger local tobacco control laws-Preemption on you	Yes			1547
Preemptive transplantation in end stage kidney failure (percent)	Percent of patients who receive a preemptive transplant at the start of ESRD	Yes			474
Pregnancies conceived within 18 months of a previous birth (percent)	Percent of pregnancies conceived within 18 months of a previous birth	Yes			784
Pregnancies: females 15-17 years (per 1,000)	Pregnancy rate among adolescent females aged 15 to 17 years	Yes			788
Pregnancies: females 18-19 years (per 1,000)	Pregnancy rate among adolescent females aged 18 to 19 years	Yes			789
Pregnancy despite use of reversible contraceptive method (percent)	Percent of females experiencing pregnancy despite use of a reversible contraceptive method	Yes			780
Prenatal care, early and adequate (percent)	Percent of pregnant women who receive early and adequate prenatal care	Yes			1132
Prenatal care, first trimester (percent)	Percent of pregnant women who receive prenatal care beginning in the first trimester of pregnancy	Yes			1131
Prescription drugs not received due to cost (percent)	Percent of persons of all ages who did not get prescription drugs in the previous 12 months due to cos	Yes	Yes		50066
Prescription pain reliever abuse (percent)	Percent of person 12 years and older with nonmedical use of prescription pain relievers in the past year	Yes			1461
Prescription psychotherapeutic drug abuse (percent)	Percent of persons 12 years and older with nonmedical use of any prescription psychotherapeutic dru	Yes			1465
Prescription sedatives abuse (percent)	Percent of persons 12 years and older with nonmedical use of prescription sedatives in the past year	Yes			1464
Prescription stimulants abuse (percent)	Percent of persons 12 years and older with nonmedical use of prescription stimulants in the past year	Yes			1463
Prescription tranquilizers abuse (percent)	Percent of persons 12 years and older with nonmedical use of prescription tranquilizers in the past year	Yes			1462
Pressure ulcer-related hospitalizations: older adults (per 100,000)	Rate of pressure ulcer-related hospitalizations among older adults	Yes			1249
Preterm births, <32 weeks of gestation (percent)	Percent of live births that are very preterm (less than 32 weeks of gestation)	Yes	Yes	Yes	1140
Preterm births, 32-33 weeks of gestation (percent)	Percent of live births at 32 to 33 weeks of gestation	Yes	Yes	Yes	1139
Preterm births, 32-36 weeks of gestation (percent)	Percent of live births that are moderately preterm (32-36 weeks gestation)	Yes	Yes	Yes	10008
Preterm births, 34-36 weeks of gestation (percent)	Percent of live births that are late preterm (34 to 36 weeks of gestation)	Yes	Yes	Yes	1138
Preterm births, total (percent)	Percent of live births that are preterm (< 37 weeks of gestation)	Yes	Yes	Yes	1137
Preventable hospitalizations: Medicare beneficiaries (per 1,000)	Discharge rate per 1,000 Medicare fee-for-service enrollees for ambulatory care sensitive conditions		Yes	Yes	27
Primary care providers (per 100,000)	Rate of primary care providers (PCP) per 100,000		Yes	Yes	25
Private health insurance: <65 years (percent) (Source: NHIS)	Percent of persons under age 65 who have private health insurance coverage	Yes			50054
Proficiency in math: 12th grade (percent)	Percent of 12th graders whose mathematics skills are at or above the proficient achievement level for	Yes			401
Proficiency in math: 4th grade (percent)	Percent of 4th graders whose mathematics skills are at or above the proficient achievement level for 4	Yes			399
Proficiency in math: 8th grade (percent)	Percent of 8th graders whose mathematics skills are at or above the proficient achievement level for 8	Yes			400
Proficiency in reading: 12th grade (percent)	Percent of 12th graders whose reading skills are at or above the proficient achievement level for 12th	Yes			398

Proficiency in reading: 4th grade (percent)	Percent of 4th graders whose reading skills are at or above the proficient achievement level for 4th grade	Yes				396
Proficiency in reading: 8th grade (percent)	Percent of 8th graders whose reading skills are at or above the proficient achievement level for 8th grade	Yes				397
Prostate cancer deaths (per 100,000)	Prostate cancer death rate	Yes				492
Prostate cancer Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have prostate cancer	Yes	Yes	Yes	Yes	305
Prostate cancer Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have prostate cancer	Yes	Yes	Yes	Yes	306
Prostate-specific antigen tests counseling: males (percent)	Percent of men who have discussed with their health care provider whether or not to have a prostate-	Yes				512
Protective eyewear use at home: adults (percent)	Percent of adults aged 18 years and over who use protective eyewear in recreational activities and ha	Yes				1573
Protective eyewear use at home: children (percent)	Percent of children and adolescents aged 6 to 17 years who use personal protective eyewear in recrea	Yes				1572
Protective gear use in intramural activities: schools (percent)	Percent of public and private schools that require students to wear appropriate protective gear when i	Yes				1083
Protective gear use in physical education: schools (percent)	Percent of public and private schools that require students to wear appropriate protective gear when i	Yes				1082
Psychiatric hospitals (number)	Number of Psychiatric hospitals participating in the Medicare program	Yes	Yes		Yes	10047
Public health majors: 4-year colleges/universities (percent)	Percent of 4-year colleges and universities that offer public health or related majors	Yes				1385
Public health minors: 4-year colleges/universities (percent)	Percent of 4-year colleges and universities that offer public health or related minors	Yes				1386
Public health system assessment: Local public health systems (percent)	Percent of State and local public health jurisdictions that conduct a public health system assessment u	Yes				1376
Public health system assessment: State public health systems (percent)	Percent of State and local public health jurisdictions that conduct a public health system assessment u	Yes				1375
Pyrethroid level: persons 6+ years (µg/g of creatinine)	Exposure to selected environmental chemicals in the population, as measured by blood and urine conc	Yes				705
Rabies surveillance, electronic: states (number)	Number of States that use electronic data from rabies animal surveillance to inform public health prev	Yes				1024
Radon mitigation systems: homes at risk for radon exposure (percent)	Percent of homes with an operating radon mitigation system for persons living in homes at risk for rad	Yes				681
Radon-reducing construction: new homes (percent)	Percent of new single family homes (SFH) constructed with radon-reducing features, especially in high	Yes				682
Read to children: parents (percent)	Percent of parents who read to their young child	Yes				740
Recent hearing exams: adults 20-69 years (percent)	Percent of adults aged 20 to 69 years who have had a hearing examination in the past 5 years	Yes				748
Recent hearing exams: older adults (percent)	Percent of adults aged 70 years and older who have had a hearing examination in the past 5 years	Yes				749
Recent hearing exams: teens (percent)	Percent of adolescents aged 12 to 19 years who have had a hearing examination in the past 5 years	Yes				750
Recess for 20+ min. in elementary school: school districts (percent)	Percent of school districts that require or recommend elementary school recess for an appropriate per	Yes				1340
Recess, required in elementary school: school districts (percent)	Percent of school districts that require regularly scheduled elementary school recess	Yes				1339
Recess, required in elementary school: States (number)	Number of States that require regularly scheduled elementary school recess	Yes				1338
Recycling of municipal waste (percent)	Percent of municipal solid waste that is recycled or composted	Yes				662
Reduced joint mobility: persons with hemophilia (percent)	Percent of persons with hemophilia who develop reduced joint mobility due to bleeding into joints	Yes				444
Reducing fat/calories in diet: adults with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report reducing the amount of fat	Yes				564
Refrigerate promptly, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Chill: Refrigerate promptly	Yes				838
Rehabilitation Hospitals (number)	Number of Rehabilitation hospitals participating in the Medicare program	Yes	Yes		Yes	10048
Religious Nonmedical Health Care Institutions (number)	Number of religious nonmedical health care institutions participating in the Medicare program	Yes	Yes		Yes	10049
Renal evaluation after kidney injury: Medicare beneficiaries (percent)	Percent of hospital patients who incurred acute kidney injury and have followup renal evaluation in 6	Yes				482
Renal function knowledge: adults aged 45+ with chronic kidney disease (percent)	Percent of adults 45 years and older with chronic kidney disease (CKD) who know they have impaired	Yes				485
Reports of fetal death, use of latest standard form: states (number)	Number of States that record vital events using the latest U.S. standard certificates and report - States	Yes				1384
Reproductive health services receipt: sexually active females (percent)	Percent of sexually active females aged 15 to 44 years who received reproductive health services	Yes				786
Reproductive health services receipt: sexually active males (percent)	Percent of sexually active males aged 15 to 44 years who received reproductive health services	Yes				787
Residential fire deaths (per 100,000)	Residential fire death rate	Yes				1050
Riding with a driver who has been drinking: teens (percent)	Percent of adolescents who report that they rode, during the previous 30 days, with a driver who had	Yes	Yes			1427
Risk perception, binge drinking: at risk teens (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Cons	Yes				1450
Risk perception, cocaine: at risk teens (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Usin	Yes				1452
Risk perception, marijuana: at risk teens (percent)	Percent of adolescents aged 12 to 17 years perceiving great risk associated with substance abuse-Smo	Yes				1451
Rotavirus vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-2 or more doses rota	Yes	Yes			1002
Rubella, U.S.-acquired (number)	Number of rubella cases (U.S.-acquired cases)	Yes				986
Rural Health Clinics (number)	Number of rural health clinics participating in the Medicare program	Yes	Yes		Yes	10050
Safe drinking water: persons served by community water (percent)	Percent of persons served by community water systems who receive a supply of drinking water that m	Yes				659
Safety at school: perception of teens' parents (percent)	Percent of adolescents whose parents consider them to be safe at school	Yes				407
Safety belt use (percent)	Percent of persons observed using safety belts	Yes				1047

Salmonella species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Reduction in infecti	Yes				825
Salmonellosis (number)	Number of reported cases of salmonellosis	Yes	Yes	Yes		10026
Saturated fat consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from saturated fat for persons aged 2 years and over	Yes				1221
School absenteeism from illness/ injury: teens aged 12-17 (percent)	Percent of adolescents missing school 11 or more days due to illness or injury	Yes				403
School athletic facilities, public access (percent)	Percent of the Nation's public and private schools that provide access to their physical activity spaces	Yes				1318
School nurse-to-student ratio: elementary schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered sci	Yes				588
School nurse-to-student ratio: high schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered sci	Yes				586
School nurse-to-student ratio: middle schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered sci	Yes				587
School nurse-to-student ratio: schools (percent)	Percent of the Nation's elementary, middle, and senior high schools that have a full-time registered sci	Yes				585
School work meaningful: teens aged 12-17 (percent)	Percent of adolescents who consider their school work to be meaningful and important	Yes				402
Schools located near highways (percent)	Number of new schools sited within 500 feet of an interstate or Federal or State highway	Yes				725
Screen time, computer: children 2-5 years (percent)	Percent of children aged 2 to 5 years who use a computer or play computer games outside of school (f	Yes				1330
Screen time, computer: children 6-14 years (percent)	Percent of children aged 6 to 14 years who use a computer or play computer games outside of school	Yes				1331
Screen time, computer: teens (percent)	Percent of adolescents in grades 9 through 12 who use a computer or play computer games outside of	Yes	Yes			1332
Screen time: children 0-2 years (percent)	Percent of children aged 0 to 2 years who view no television or videos on an average weekday	Yes				1326
Screen time: children 2-5 years (percent)	Percent of children aged 2 to 5 years who view television, videos, or play video games for no more tha	Yes				1327
Screen time: children 6-14 years (percent)	Percent of children aged 6 to 14 years who view television, videos, or play video games for no more th	Yes				1328
Screen time: teens (percent)	Percent of adolescents in grades 9 through 12 who view television, videos, or play video games for no	Yes	Yes			1329
Secondhand smoke exposure: adult nonsmokers (percent)	Percent of nonsmokers exposed to secondhand smoke-Adults aged 18 years and older	Yes				1519
Secondhand smoke exposure: nonsmokers 12-17 years (percent)	Percent of nonsmokers exposed to secondhand smoke-Adolescents aged 12 to 17 years	Yes				1518
Secondhand smoke exposure: nonsmokers 3-11 years (percent)	Percent of nonsmokers exposed to secondhand smoke-Children aged 3 to 11 years	Yes				1517
Self-harm injuries, nonfatal (per 100,000)	Nonfatal intentional self-harm injury rate	Yes				1100
Separate food, food safety practice (percent)	Percent of consumers who follow key food safety practices-- Separate: Don't cross-contaminate	Yes				836
Septicemia deaths (per 100,000)	Septicemia death rate	Yes	Yes	Yes		50069
Serious emotional/behavioral difficulties: children (percent)	Percent of children under age 18 years with serious emotional or behavioral difficulties	Yes				50034
Serious property crimes with perpetrators 10-24 years (per 100,000)	Rate of minor and young adult perpetration of serious property crimes	Yes				414
Serious psychological distress: adults (percent)	Percent of adults 18 years of age and over who have experienced serious psychological distress in the	Yes				50055
Serious psychological distress: adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who report serious psychological distress	Yes				435
Serious violent incidents in public schools (percent)	Percent of public schools with a serious violent incident	Yes				408
Shigellosis (number)	Number of reported cases of shigellosis	Yes	Yes	Yes		10027
Shingles vaccination: older adults (percent)	Percent of adults who are vaccinated against zoster (shingles)	Yes				1027
Short stay hospitals (number)	Number of short stay hospitals participating in the Medicare program	Yes	Yes		Yes	10051
Sick leave use: adults with balance/dizziness problems (percent)	Percent of adults who have missed days of work or school in the past 12 months because of balance ar	Yes				768
Single-parent households (percent)	Percent of households that are single-parent households				Yes	40
Skilled nursing facility Medicare admissions (per 1,000 beneficiaries)	Rate of skilled nursing facility (SNF) admissions among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2008
Skilled nursing facility Medicare days (per 1,000 beneficiaries)	Average number of skilled nursing facility (SNF) days among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	2016
Skilled nursing facility Medicare users (count)	Medicare fee-for-service beneficiaries who had covered skilled nursing facility services	Yes	Yes	Yes	Yes	214
Skilled nursing facility Medicare utilization (percent)	Medicare fee-for-service beneficiaries who had covered skilled nursing facility services	Yes	Yes	Yes	Yes	215
Skilled Nursing Home Title 18 Facilities (number)	Number of skilled nursing home Title 18 (Medicare only) facilities	Yes	Yes		Yes	10052
Skin allergy: children (percent)	Percent of children under age 18 years who have a skin allergy	Yes				50036
Skin cancer protective measures: adults (percent)	Percent of adults aged 18 years and older who follow protective measures that may reduce the risk of	Yes				504
Skin cancer protective measures: teens (percent)	Percent of adolescents in grades 9 through 12 who follow protective measures that may reduce the ris	Yes				503
Skin diseases or disorders, work-related (per 10,000)	Rate of occupational skin diseases or disorders among full-time workers	Yes				1301
Smoke-free air laws covering bars: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes				1527
Smoke-free air laws covering commercial daycare centers: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes				1529
Smoke-free air laws covering home based daycare centers: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes				1530
Smoke-free air laws covering hotels and motels: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes				1532
Smoke-free air laws covering multiunit housing: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes				1533

Smoke-free air laws covering Prisons: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1535
Smoke-free air laws covering private workplaces: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1524
Smoke-free air laws covering public transportation: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1531
Smoke-free air laws covering public workplaces: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1525
Smoke-free air laws covering restaurants: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1526
Smoke-free air laws covering vehicles with children: States (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1534
Smoke-free homes: adults (percent)	Percent of adults living in smoke-free homes	Yes			1561
Smoke-free indoor air laws--college/university campuses (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1540
Smoke-free indoor air laws--entrances/exits of all public places (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1538
Smoke-free indoor air laws--Gaming Halls (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1528
Smoke-free indoor air laws--Hospital campuses (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1539
Smoke-free indoor air laws--mental health treatment facilities (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1537
Smoke-free indoor air laws--Substance abuse treatment facilities (number)	Number of States, District of Columbia, Territories, and Tribes with established laws on smoke-free inc	Yes			1536
Smokeless tobacco use in past month: teens (percent)	Percent of adolescent reporting use of smokeless tobacco products in the past month - Students (grad	Yes	Yes		1503
Smokeless tobacco use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating smokeless tobacco use in the past 12 months	Yes			1511
Smokeless tobacco use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating smokeless tobacco use in past 12 mc	Yes			1507
Smokeless tobacco use: adults (percent)	Percent of adults aged 18 years and over using smokeless tobacco products	Yes			1499
Smoking abstinence, preconception (percent)	Percent of women delivering a live birth who did not smoke prior to pregnancy	Yes			1180
Smoking cessation attempts: adult smokers (percent)	Percent of adult smokers aged 18 years and over who attempted smoking cessation in the past 12 mo	Yes			1513
Smoking cessation attempts: teen smokers (percent)	Percent of adolescent smokers in grades 9 through 12 who attempted smoking cessation in the past 1:	Yes	Yes		1515
Smoking cessation during pregnancy (percent)	Percent of females aged 18 to 49 years who ceased smoking during the first trimester of pregnancy	Yes			1491
Smoking cessation success (percent)	Percent of adults 18 years and older with smoking cessation success--6 months to 1 year ago	Yes			1555
Social and emotional support: adults with disabilities (percent)	Percent of adults with disabilities who report sufficient social and emotional support	Yes	Yes		515
Social support for health concerns (percent)	Percent of adults who report having friends or family members whom they talk with about their healt	Yes			870
Social-emotional support lacking: Adults (percent)	Percent of adults 18 years and over who report not receiving sufficient social-emotional support		Yes	Yes	50028
Sodium consumption (milligrams)	Mean total daily sodium intake (in mg) by persons aged 2 years and over	Yes			1222
Solid fat and added sugar consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from solid fats and added sugars for persons aged 2 years ar	Yes			1238
Solid fat consumption (mean percentage of total daily calorie intake)	Mean percent of total daily calorie intake from solid fats for persons aged 2 years and over	Yes			1236
Source of ongoing care: <18 years (percent)	Percent of persons who have a specific source of ongoing care--children and youth aged 17 years and	Yes			376
Source of ongoing care: 18 to 64 years (percent)	Percent of persons who have a specific source of ongoing care--adults aged 18 to 64 years	Yes			377
Source of ongoing care: 65+ years (percent)	Percent of persons who have a specific source of care - adults aged 65 years and older	Yes			378
Source of ongoing care: all ages (percent)	Percent of persons who have a specific source of ongoing care--all ages	Yes			375
Source of usual care, none: adults <65 years (percent)	Percent of adults aged 18-64 with no usual source of health care	Yes			50061
Source of usual care, none: children (percent)	Percent of children under age 18 years who have no usual source of health care	Yes			50060
Special services: children with Autism spectrum disorders (percent)	Percent of children with an ASD enrolled in special services by 48 months of age	Yes			1190
Specialty health care use: adults with balance/dizziness problems (percent)	Percent of adults with moderate to severe balance or dizziness problems who have seen or been refer	Yes			763
Specialty health care use: adults with tinnitus (percent)	Percent of adults bothered by the onset of tinnitus in the past 5 years, for whom it is a moderate, big,	Yes			760
Spina bifida (per 100,000 live births)	Rate of spina bifida	Yes			1169
Spinal cord injury deaths (per 100,000)	Spinal cord injury death rate	Yes	Yes		1062
Spinal cord injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal traumatic spinal cord injuries	Yes			1063
Spotted Fever Rickettsiosis (number)	Number of reported cases of spotted fever rickettsiosis	Yes	Yes	Yes	10028
State epidemiologists with formal training in epidemiology (percent)	Percent of Tribal, State, and local public health agencies that provide or assure comprehensive epidem	Yes			1360
Statewide cancer registries (number)	Number of central, population-based registries from the 50 States and the District of Columbia that ca	Yes			497
STD discussion with parent before age 18: females (percent)	Percent of adolescents females who talked to a parent or guardian about STDs before they were 18 ye	Yes			816
STD discussion with parent before age 18: males (percent)	Percent of adolescents males who talked to a parent or guardian about STDs before they were 18 year	Yes			817
STD instruction receipt before age 18: females (percent)	Percent of adolescents females who received formal instruction on STDs before they were 18 years olc	Yes			808
STD instruction receipt before age 18: males (percent)	Percent of adolescents males who received formal instruction on STDs before they were 18 years old	Yes			809
Steps to reduce risk promoted: news stories on foodborne disease outbreaks or national disasters (per	Percent of crisis and emergency risk messages embedded in print and broadcast news stories that pro	Yes			10097

Steroid use: 10th graders (percent)	Percent of 10th graders using steroids	Yes				1441
Steroid use: 8th graders (percent)	Percent of 8th graders using steroids	Yes				1440
Steroid use: high school seniors (percent)	Percent of 12th graders using steroids	Yes				1442
Streptococcus pneumoniae, drug-resistant (number)	Number of reported cases of drug-resistant streptococcus pneumoniae	Yes	Yes	Yes		10025
Stroke awareness: importance of calling 9-1-1 (percent)	Percent of adults aged 20 years and over who are aware of the importance of accessing rapid emerger	Yes				889
Stroke awareness: symptoms (percent)	Percent of adults aged 20 years and over who are aware of the early warning symptoms and signs of a	Yes				888
Stroke awareness: symptoms and importance of calling 9-1-1 (percent)	Percent of adults who are aware of the early warning symptoms and signs of a stroke and the importa	Yes				887
Stroke deaths (per 100,000)	Stroke death rate	Yes	Yes	Yes		881
Stroke Medicare beneficiaries (count)	Medicare fee-for-service beneficiaries who have had a stroke	Yes	Yes	Yes	Yes	50020
Stroke Medicare beneficiaries (percent)	Medicare fee-for-service beneficiaries who have had a stroke	Yes	Yes	Yes	Yes	50025
Stroke prevalence: adults (percent)	Respondent-reported prevalence of stroke among adults aged 18 and over	Yes				50046
Substance abuse prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services,	Yes				600
Sufficient sleep: adults (percent)	Percent of adults who get sufficient sleep	Yes				1472
Sufficient sleep: teens (percent)	Percent of students in grades 9 through 12 who get sufficient sleep	Yes				1471
Suffocation deaths, unintentional (per 100,000)	Unintentional suffocation death rate: All persons	Yes				1068
Suffocation deaths, unintentional: infants (per 100,000)	Unintentional suffocation death rate: Infants 0 to 12 months	Yes				1069
Suffocation deaths, unintentional: older adults (per 100,000)	Unintentional suffocation death rate: Adults aged 65 years and over	Yes				1070
Suicide attempts requiring medical attention: teens (per 100)	Rate of suicide attempts by adolescents that require medical treatment	Yes	Yes			1106
Suicide deaths (per 100,000)	Suicide death rate	Yes	Yes	Yes		1105
Sunburn: adults (percent)	Percent of adults aged 18 years and older who report sunburn	Yes				500
Surgery patients who received appropriate venous thromboembolism prophylaxis (percent)	Hospital Compare: Surgery patients who received appropriate venous thromboembolism prophylaxis	Yes	Yes	Yes	Yes	342
Surgery patients with appropriate hair removal (percent)	Hospital Compare: Surgery patients with appropriate hair removal	Yes	Yes	Yes	Yes	340
Surgery patients with appropriate prophylactic antibiotic selection (percent)	Hospital Compare: Surgery patients with appropriate prophylactic antibiotic selection	Yes	Yes	Yes	Yes	337
Surgery patients with prophylactic antibiotic received within one hour prior to surgery incision (percen	Hospital Compare: Surgery patients with prophylactic antibiotic received within one hour prior to surg	Yes	Yes	Yes	Yes	336
Surgery patients with prophylactic antibiotics discontinued within 24 hours after surgery end time (per	Hospital Compare: Surgery patients with prophylactic antibiotics discontinued within 24 hours after su	Yes	Yes	Yes	Yes	338
Surgery patients with recommended venous thromboembolism prophylaxis ordered (percent)	Hospital Compare: Surgery patients with recommended venous thromboembolism prophylaxis ordere	Yes	Yes	Yes	Yes	341
Surveillance programs for caregivers: States (number)	States and the District of Columbia with surveillance programs for caregivers	Yes				528
Survival after AIDS diagnosis (percent)	Percent of persons surviving more than 3 years after a diagnosis with AIDS	Yes				937
Sweetened beverages not offered: schools (percent)	Percent of schools that do not sell or offer calorically sweetened beverages to students.	Yes				10065
Syphilis, congenital (per 100,000)	Congenital syphilis rate	Yes				1474
Syphilis, primary and secondary: females (per 100,000)	Domestic transmission of primary and secondary syphilis among females	Yes				1480
Syphilis, primary and secondary: males (per 100,000)	Domestic transmission of primary and secondary syphilis among males	Yes				1481
TB curative therapy completion (percent)	Percent of all tuberculosis patients who complete curative therapy within 12 months	Yes				970
TB test confirmation timeliness (percent)	Average time for a laboratory to confirm and report tuberculosis cases	Yes				972
TB treatment: persons with latent TB (percent)	Percent of persons with latent tuberculosis infection who complete a course of treatment	Yes				971
Tdap vaccination: teens 13-15 years (percent)	Percent of adolescents who have 1 dose of tetanus-diphtheria-acellular pertussis (Tdap) booster vaccir	Yes	Yes			1010
Telecommuting (percent)	Percent of trips to work using of alternative modes of transportation - telecommute	Yes				669
Test Medicare service events fee-for-service beneficiaries (per 1,000 beneficiaries)	Rate of test service events among Medicare fee-for-service beneficiaries	Yes	Yes	Yes	Yes	50072
Test Medicare users (count)	Medicare fee-for-service beneficiaries who used test services	Yes	Yes	Yes	Yes	50070
Test Medicare utilization (percent)	Medicare fee-for-service beneficiaries who used test services	Yes	Yes	Yes	Yes	50071
Timely evaluation: children with Autism spectrum disorder (percent)	Percent of children with an ASD with a first evaluation by 36 months of age	Yes				1189
Tobacco cessation counseling in dental care settings: persons 12+ years (percent)	Percent of tobacco users ages 12 and older who receive tobacco cessation counseling at dental visits	Yes				1553
Tobacco cessation counseling in hospital outpatient dept. visits: adults (percent)	Percent of tobacco users ages 18 and older who receive tobacco cessation counseling at hospital ambu	Yes				1552
Tobacco cessation counseling in physician office visits: adults (percent)	Percent of tobacco users ages 18 and older who receive tobacco cessation counseling at office based a	Yes				1551
Tobacco free environments: high schools (percent)	Percent of senior high schools with tobacco-free environments	Yes				1522
Tobacco free environments: junior high schools (percent)	Percent of Junior high schools with tobacco-free environments	Yes				1520
Tobacco free environments: middle schools (percent)	Percent of middle schools with tobacco-free environments	Yes				1521
Tobacco marketing exposure, internet: teens (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco marketing-I	Yes				1541

Tobacco marketing exposure, magazine/newspaper: teens (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco marketing-I	Yes				1542
Tobacco marketing exposure, point of purchase: teens (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco marketing a	Yes				1544
Tobacco marketing exposure, television/movies: teens (percent)	Percent of adolescents and young adults grades 6 through 12 who are exposed to tobacco marketing i	Yes				1543
Tobacco screening in dental care setting: persons 12+ years (percent)	Percent of persons aged 12 years and older who received tobacco screening in dental care settings	Yes				1559
Tobacco screening in hospital outpatient dept. visits: adults (percent)	Percent of persons aged 18 years and older who received tobacco screening in hospital ambulatory ca	Yes				1558
Tobacco screening in physician office visits: adults (percent)	Percent of persons aged 18 years and older who received tobacco screening in office-based ambulator	Yes				1557
Tobacco tax increases on cigarettes: States/Federal (number)	Number of States, DC, and Federal government that have increased tax on cigarettes by \$1.50	Yes				1548
Tobacco tax increases on smokeless tobacco: States/Federal (number)	Number of States, DC, and Federal government that have increased tax on smokeless tobacco product	Yes				1549
Tobacco use in past month: teens (percent)	Percent of adolescent reporting use of tobacco products in the past month - Students (grades 9 throug	Yes	Yes			1501
Tobacco use initiation: adults 18-25 years (percent)	Percent of young adults aged 18 to 25 years initiating tobacco products use in the past 12 months	Yes				1509
Tobacco use initiation: teens (percent)	Percent of children and adolescents aged 12 to 17 years initiating tobacco use in past 12 months	Yes				1505
Tobacco use prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services,	Yes				599
Tooth loss, all permanent: adults 65-74 years (percent)	Percent of older adults aged 65 to 74 years who have lost all of their natural teeth	Yes				1279
Tooth loss, any permanent: adults 45-64 years (percent)	Percent of adults aged 45 to 64 years who have ever had a permanent tooth extracted because of den	Yes				1278
Toxic chemicals (pounds)	Toxic chemicals released annually	Yes	Yes	Yes		105
Toxic pollutants (tons)	Amount of toxic pollutants released into the environment	Yes				678
Transition planning: teens with special health care needs (percent)	Percent of youth with special health care needs whose health care provider has discussed transition pl	Yes				536
Transplant Centers (number)	Number of transplant centers	Yes	Yes	Yes		50026
Trauma care access, by land mass (percent)	Percent of the land mass of the continental United States with access to trauma care	Yes				1097
Trauma care access, by population (percent)	Percent of the population residing within the continental United States with access to trauma care	Yes				1096
Traumatic brain injury deaths (per 100,000)	Traumatic brain injury death rate	Yes	Yes			1059
Traumatic brain injury ED visits (per 100,000)	Rate of emergency department visits for nonfatal traumatic brain injuries	Yes				1061
Traumatic brain injury hospitalizations (per 100,000)	Hospitalization rate for nonfatal traumatic brain injuries	Yes				1060
Treatment: adults with both substance abuse and mental disorders (percent)	Percent of persons with co-occurring substance abuse and mental disorders who receive treatment fo	Yes				1119
Treatment: adults with major depressive episode (percent)	Percent of adults aged 18 years and older with major depressive episode	Yes				1118
Treatment: adults with serious mental illness (percent)	Percent of adults aged 18 years and older with serious mental illness (SMI)	Yes				1117
Treatment: children with mental health problems (percent)	Percent of children with mental health problems who receive treatment	Yes				1110
Trying to lose weight: adults with pre-diabetes (percent)	Percent of persons at high risk for diabetes (with pre-diabetes) who report trying to lose weight	Yes				563
Tuberculosis (per 100,000)	Tuberculosis (TB) rate	Yes				969
Tuberculosis: foreign born persons (per 100,000)	Tuberculosis (TB) case rate for foreign-born persons living in the United States	Yes				845
Unemployed: persons 16+ (percent)	Percent of population age 16+ unemployed and looking for work	Yes	Yes	Yes		35
Unemployment: adults 18-64 years with arthritis (percent)	Unemployment rate among adults aged 18-64 years with doctor-diagnosed arthritis	Yes				423
Unemployment: persons 16-64 years with disabilities (percent)	Unemployment rates among adults aged 16-64 with disabilities	Yes				541
Unintended pregnancy prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services,	Yes				601
Unintentional Injury deaths: No MVA (per 100,000)	Unintentional injury death rate (excluding motor vehicle injuries)	Yes	Yes	Yes		79
Unknown risks to health presented: news stories on foodborne disease outbreaks or national disasters	Percent of crisis and emergency risk messages embedded in print and broadcast news stories that expl	Yes				10095
Urinary microalbumin measurement: diabetic Medicare beneficiaries (percent)	Percent of persons with diagnosed diabetes who obtain an annual urinary microalbumin measurement	Yes				549
Urinary tract infection admission rate (per 100,000 beneficiaries)	Prevention Quality Indicators: Urinary tract infection admission rate among Medicare fee-for-service b	Yes	Yes	Yes	Yes	2044
Use of hearing protection: adults (per 1,000)	Rate of adults aged 20 to 69 years who have ever used hearing protection devices (earplugs, earmuffs)	Yes				752
Use of hearing protection: teens (per 1,000)	Rate of adolescents aged 12 to 19 years who have ever used hearing protection devices (earplugs, ear	Yes				753
Use of mobile devices for internet access (percent)	Percent of persons who use mobile devices for internet use	Yes				859
Usual primary care provider (percent)	Percent of persons with a usual primary care provider	Yes				372
Uterine cervix cancer deaths (per 100,000)	Uterine cervix cancer death rate	Yes				489
Vaccination coverage assessment: private health providers (percent)	Percent of private providers who have had vaccination coverage levels among children in their practic	Yes				977
Vaccination coverage assessment: public health providers (percent)	Percent of public health providers who have had vaccination coverage levels among children in their p	Yes				976
Vaccination record keeping (percent)	Percent of children under age 6 years of age whose immunization records are in fully operational, pop	Yes				1009
Vaccination, zero doses: children 18-35 months (percent)	Percent of children in the United States who receive zero doses of recommended vaccines by age 19 to	Yes				1039
Varicella vaccination: children 19-35 months (percent)	Percent of young children who have received universally recommended vaccines-1 dose varicella vacci	Yes	Yes			998

Varicella vaccination: kindergarteners (percent)	Vaccination coverage levels for children in kindergarten-Varicella vaccine (2 or more doses)	Yes			1007
Varicella vaccination: teens 13-15 years (percent)	Percent of adolescents who have 2 doses of varicella vaccine by age 13 to 15 years (excluding children	Yes	Yes		1011
Varicella: children (number)	Number of varicella (chicken pox) cases among persons aged 17 years of age and under	Yes			987
Vascular access for hemodialysis (percent)	Percent of adult hemodialysis patients who use arteriovenous fistulas as the primary mode of vascular	Yes			469
Vegetable consumption (cup equivalents per 1,000 calories)	Mean daily intake of vegetables (in cup equivalents per 1,000 calories) by persons aged 2 years and ov	Yes			1218
Vegetable consumption-dark green, orange, and legumes (cup equivalents per 1,000 calories)	Mean daily intake of dark green vegetables, orange vegetables, and legumes (in cup equivalents per 1,	Yes			1219
Vehicle restraint proper use: children 1-3 years (percent)	Percent of age-appropriate vehicle restraint system use in children aged 1 to 3 years	Yes			1055
Vehicle restraint proper use: children 4-7 years (percent)	Percent of age-appropriate vehicle restraint system use in children aged 4 to 7 years	Yes			1056
Vehicle restraint proper use: children 8-12 years (percent)	Percent of age-appropriate vehicle restraint system use in children aged 8 to 12 years	Yes			1057
Vehicle restraint proper use: infants 0-12 months (percent)	Percent of age-appropriate vehicle restraint system use in infants 0 to 12 months	Yes			1054
Venous thromboembolism physician visits (per 10,000)	Rate of visits for venous thromboembolism (VTE)	Yes			441
Very low birth weight deliveries (percent)	Percent of live births that are very low birth weight (VLBW)	Yes	Yes	Yes	1136
Very low birth weight infants born at level III hospitals (percent)	Percent of very low birth weight (VLBW) infants born at level III hospitals or subspecialty perinatal cen	Yes			1195
Vibrio species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Vibrio species	Yes			827
Violence exposure: children (percent)	Percent of children who are exposed to violence	Yes			1104
Violence prevention: community-based organizations (percent)	Percent of community-based organizations (including local health departments, tribal health services	Yes			597
Violent crimes: perpetrators 10-24 years (per 100,000)	Rate of minor and young adult perpetration of violent crimes	Yes			413
Violent death surveillance: states (number)	Number of states that link data on violent deaths from death certificates, law enforcement, and coron	Yes			1101
Vision limitations: adults (percent)	Percent of adults aged 18 and over with any trouble seeing, even with glasses or contacts	Yes			50047
Vision screening: children & 5 years (percent)	Percent of preschool children aged 5 years and under who receive vision screening	Yes			1562
Visitable features: homes and residential buildings (percent)	Percent of newly constructed or retrofitted U.S. homes and residential buildings that have visitable fea	Yes			542
Visual impairment due to cataracts: older adults (per 1,000)	Rate of visual impairment due to cataract	Yes			1570
Visual impairment due to diabetic retinopathy: adults with diabetes (per 1,000)	Rate of visual impairment due to diabetic retinopathy	Yes			1568
Visual impairment due to macular degeneration: adults 45+ years (per 1,000)	Rate of visual impairment due to age-related macular degeneration (AMD)	Yes			1571
Visual impairment due to refractive errors: persons 12+ years (per 1,000)	Rate of visual impairment due to uncorrected refractive error	Yes			1567
Visual impairment: children <18 years (per 1,000)	Percent of children and adolescents aged 17 years and under with blindness and visual impairment	Yes			1563
Visual impairments due to glaucoma: adults 45+ years (per 1,000)	Rate of visual impairment due to glaucoma	Yes			1569
Visual rehabilitation services use: adults with visual impairments (per 1,000)	Rate of use of vision rehabilitation services by persons with visual impairment	Yes			1574
von Willebrand Disease diagnosis: women (percent)	Percent of women with von Willebrand disease (vWD) who are timely and accurately diagnosed	Yes			443
Walking for transportation (percent)	Percent of trips to work using of alternative modes of transportation - walking	Yes			667
Water quality/sanitation/hygiene deaths: global (number in thousands)	Global burden of disease resulting in death due to poor water quality, sanitation, and insufficient hygie	Yes			720
Waterborne disease outbreaks (number)	Number of waterborne disease outbreaks arising from water intended for drinking among persons ser	Yes			660
Weapon carrying on school property: teens (percent)	Percent of weapon carrying by adolescents on school property	Yes	Yes		1053
Weight loss counseling: overweight adults with arthritis (percent)	Percent of adults with doctor-diagnosed arthritis who receive health care provider counseling-For wei	Yes			421
Weight loss, nutrition or physical activity counseling: obese patients (percent)	Percent of physician office visits made by adult patients who are obese that include counseling or educ	Yes			1226
Welcome to Medicare benefit use (percent)	Percent of older adults who use the Welcome to Medicare benefit	Yes			1260
Wellness check-up: teens aged 12-17 (percent)	Percent of adolescents who have had a wellness checkup in the past 12 months	Yes			405
Whole grain consumption (ounce equivalents per 1,000 calories)	Mean daily intake of whole grains (in ounce equivalents per 1,000 calories) by persons aged 2 years an	Yes			1220
Work limitations: adults 18-64 years with arthritis (percent)	Percent of adults aged 18-64 years with doctor-diagnosed arthritis who are limited in their ability to w	Yes			424
Worksite lactation support programs (percent)	Percent of employers that have worksite lactation support programs	Yes			1184
Years of potential life lost before age 75 (per 100,000)	Rate per 100,000 of years of potential life lost before age 75	Yes	Yes	Yes	3
Yersinia species infections (per 100,000)	Rate of infections caused by key pathogens transmitted commonly through food: Yersinia species	Yes			828